Please check the examination details be	low before ente	ering your candidate information
Candidate surname		Other names
Centre Number Candidate N	umber	
Pearson Edexcel Inter	nation	al Advanced Level
Time 1 hour 30 minutes	Paper reference	WMA13/01
Mathematics		
International Advanced L Pure Mathematics P3	evel	
You must have: Mathematical Formulae and Statistic	al Tables (Ye	Pllow), calculator

Candidates may use any calculator permitted by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

$$f(x) = \frac{2x^3 - 4x - 15}{x^2 + 3x + 4}$$

(a) Show that

$$f(x) \equiv Ax + B + \frac{C(2x+3)}{x^2 + 3x + 4}$$

where A, B and C are integers to be found.


(4)

(5)

(b) Hence, find

$$\int_{3}^{5} f(x) dx$$

giving your answer in the form $p + \ln q$, where p and q are integers.



	Le bl
nestion 1 continued	



nestion 1 continued	

Question 1 continued	blank
(Total 9 marks)	Q1
, , ,	



2. The functions f and g are defined by

$$f(x) = 5 - \frac{4}{3x + 2} \qquad x \geqslant 0$$

$$g(x) = \left| 4\sin\left(\frac{x}{3} + \frac{\pi}{6}\right) \right| \quad x \in \mathbb{R}$$

(a) Find the range of f

(2)

- (b) (i) Find $f^{-1}(x)$
 - (ii) Write down the domain of f^{-1}

(3)

(c) Find $fg(-\pi)$

(2)



Question 2 continued	blank
	Q2
(Total 7 marks)	



In this question you must show all stages of your working.
 Solutions relying entirely on calculator technology are not acceptable.




Figure 1

Figure 1 shows a sketch of part of the curve with equation y = f(x) where

$$f(x) = (x-2)^2 e^{3x} \qquad x \in \mathbb{R}$$

The curve has a maximum turning point at A and a minimum turning point at (2, 0)

(a) Use calculus to find the exact coordinates of A.

(5)

Given that the equation f(x) = k, where k is a constant, has at least two distinct roots,

(b) state the range of possible values for k.

(2)

Question 3 continued	blank
	Q3
(Total 7 marks)	



4. $y = \log_{10}(2x + 1)$	
(a) Express x in terms of y .	(2)
(b) Hence, giving your answer in terms of x , find $\frac{dy}{dx}$	(3)

Question 4 continued		blank
		Q4
	(Total 5 marks)	



5.




Figure 2

The profit made by a company, $\pounds P$ million, t years after the company started trading, is modelled by the equation

$$P = \frac{4t - 1}{10} + \frac{3}{4} \ln \left[\frac{t + 1}{(2t + 1)^2} \right]$$

The graph of P against t is shown in Figure 2.

According to the model,

(a) show that exactly one year after it started trading, the company had made a loss of approximately $\pounds\,830\,000$

(2)

A manager of the company wants to know the value of t for which P = 0

(b) Show that this value of t occurs in the interval [6, 7] (2)

(c) Show that the equation P = 0 can be expressed in the form

$$t = \frac{1}{4} + \frac{15}{8} \ln \left[\frac{(2t+1)^2}{t+1} \right]$$
 (2)

(d) Using the iteration formula

$$t_{n+1} = \frac{1}{4} + \frac{15}{8} \ln \left[\frac{\left(2t_n + 1\right)^2}{t_n + 1} \right]$$
 with $t_1 = 6$

find the value of t_2 and the value of t_6 , giving your answers to 3 decimal places.

(3)

(e) Hence find, according to the model, how many months it takes in total, from when the company started trading, for it to make a profit.

(2)

estion 5 continued	



Question 5 continued		

uestion 5 continued		Leave
	Duestion 5 continued	blank
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
(Total 11 marks)		Q5
	(Tatal 11 maul	ke)



6. $y = \frac{2 + 3\sin x}{\cos x + \sin x}$

Show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{a\tan x + b\sec x + c}{\sec x + 2\sin x}$$

where a, b and c are integers to be found.

14.	
	(6)

	Le bl
Question 6 continued	l bi



Question 6 continued	Leave
Auction a continuen	
	Q6
(Total 6 marks)	






Figure 3

Figure 3 shows a sketch of the graph of C_1 with equation

$$y = 5 - |3x - 22|$$

- (a) Write down the coordinates of
 - (i) the vertex of C_1
 - (ii) the intersection of C_1 with the y-axis.

(2)

(b) Find the x coordinates of the intersections of C_1 with the x-axis.

(2)

Diagram 1, shown on page 21, is a copy of Figure 3.

(c) On Diagram 1, sketch the curve C_2 with equation

$$y = \frac{1}{9}x^2 - 9$$

Identify clearly the coordinates of any points of intersection of C_2 with the coordinate axes.

(3)

(d) Find the coordinates of the points of intersection of C_1 and C_2 (Solutions relying entirely on calculator technology are not acceptable.)

(5)

Question 7 continued
<i>y</i> ♠
O X
C_1
/ \
Diagram 1

ued		

	Leave blank
Question 7 continued	biank
	0=
	Q7
(Total 12 marks)	
(1000112 1101115)	



8. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Express $8\sin x - 15\cos x$ in the form $R\sin(x-\alpha)$, where R > 0 and $0 < \alpha < \frac{\pi}{2}$

Give the exact value of R, and give the value of α , in radians, to 4 significant figures.

$$f(x) = \frac{15}{41 + 16\sin x - 30\cos x} \qquad x > 0$$

- (b) Find
 - (i) the minimum value of f(x)
 - (ii) the smallest value of x at which this minimum value occurs.

(4)

(c) State the y coordinate of the minimum points on the curve with equation

$$y = 2f(x) - 5$$
 $x > 0$ (1)

(d) State the smallest value of x at which a maximum point occurs for the curve with equation

$$y = -f(2x)$$
 $x > 0$ (1)

	b
estion 8 continued	



	Le
Question 8 continued	bla
Question o continueu	

Question 8 continued	blank
	Q8
(Total 9 marks)	



In this question you must show all stages of your working.
 Solutions relying entirely on calculator technology are not acceptable.

Given that $\cos 2\theta - \sin 3\theta \neq 0$

(a) prove that

$$\frac{\cos^2 \theta}{\cos 2\theta - \sin 3\theta} \equiv \frac{1 + \sin \theta}{1 - 2\sin \theta - 4\sin^2 \theta}$$

(4)

(b) Hence solve, for $0 < \theta \leqslant 360^{\circ}$

$$\frac{\cos^2 \theta}{\cos 2\theta - \sin 3\theta} = 2 \csc \theta$$

Give your answers to one decimal place.

(5)



	Le
nestion 9 continued	bl

	Le
nestion 9 continued	bl

()	9

(Total 9 marks)

TOTAL FOR PAPER IS 75 MARKS

END



Question 9 continued