Instructions for the Examinations Officer

Pearson Edexcel International A level Mathematics Mechanics Paper 1 WME01 Date: 10 January 2025

There is an error on page 2 of this question paper. A Clarification Notice is enclosed below.

At the start of the examination the Clarification Notice must be read aloud to the candidates taking the examination.

All other questions on the paper have been printed correctly.

Clarification Notice

There is an error on page 2, Question 1b.

Please amend your question paper as follows:

The question currently reads:

- The resultant of the four forces acts in **the direction of** the vector (7i +2j). It should read:
- The resultant of the four forces acts in a direction that is parallel to the vector (7i + 2j).

Please check the examination details belo	w before ente	ring your candidate information
Candidate surname		Other names
Centre Number Candidate Nu	mber	
Pearson Edexcel International Advanced Level		
Friday 10 January 2025		
Afternoon (Time: 1 hour 30 minutes)	Paper reference	WME01/01
Mathematics		
	hcidiar	v/Advancod Lovol
International Advanced Subsidiary/Advanced Level		
Mechanics M1		
You must have: Total Marks		
Mathematical Formulae and Statistical Tables (Yellow), calculator		

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided

 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over







1. A particle of mass 2.5 kg moves on a smooth horizontal plane under the action of three horizontal forces, \mathbf{F}_1 , \mathbf{F}_2 and \mathbf{F}_3 , where

$$\mathbf{F}_1 = (6\mathbf{i} + 8\mathbf{j})\,\mathbf{N}$$

$$\mathbf{F}_2 = (-16\mathbf{i} + 2\mathbf{j})\,\mathbf{N}$$

$$\mathbf{F}_3 = (-2\mathbf{i} + 8\mathbf{j})\,\mathbf{N}$$

(a) Find the magnitude of the acceleration of the particle.

(4)

A fourth force, $\mathbf{F}_4 = (p\mathbf{i} + p\mathbf{j})\mathbf{N}$, where p is a constant, is added.

The resultant of the four forces acts in the direction of the vector $(7\mathbf{i} + 2\mathbf{j})$.

(b) Find the value of *p*.

(4)

Question 1 continued	
(T.	otal for Question 1 is 8 marks)
	om ioi Question i is o marks)



- 2. The fixed points A, B and C lie in a straight line on a horizontal road.
 - At time t = 0, a motorbike passes through A with speed $5 \,\mathrm{m \, s}^{-1}$
 - From A, the motorbike accelerates uniformly until it reaches B with a speed of $V \text{m s}^{-1}$
 - The motorbike takes T_1 seconds to travel from A to B
 - From B, the motorbike decelerates uniformly until it comes to rest at C
 - The motorbike takes T_2 seconds to travel from B to C
 - (a) Sketch a speed-time graph for the motion of the motorbike as it moves from A to C.

(3)

The distance AB is 132 m and the distance BC is 136 m.

- (b) Find, in terms of V, an expression for
 - (i) T_1
 - (ii) T_2

(4)

Given that the motorbike takes 28 s to travel from A to C,

(c) find the value of V,

(2)

(d) find the deceleration of the motorbike.

(2)

Question 2 continued		



Question 2 continued

Question 2 continued	
(Total	for Questian 2 is 11 mayles)
(10tal	for Question 2 is 11 marks)



3.




Figure 1

A particle P of mass $3m \log$ and a particle Q of mass $m \log$ are at rest on a rough horizontal surface. The distance between P and Q is 1.4m, as shown in Figure 1.

An impulse of magnitude λNs is now applied to P in the direction PQ. Immediately after the impulse is applied, the speed of P is 5 m s^{-1}

(a) Find λ in terms of m.

(2)

Immediately before P collides with Q, the speed of P is $2.5\,\mathrm{m\,s}^{-1}$ The coefficient of friction between P and the surface is μ

(b) Find the value of μ

(7)

Immediately after P collides with Q, the speed of Q is $2.1 \,\mathrm{m\,s}^{-1}$

(c) Find the speed of P immediately after the collision.

(3)



Question 3 continued		



Question 3 continued		

Question 3 continued	
(Total	al for Question 3 is 12 marks)
(100)	() () () () () () () () () ()



4.




Figure 2

A uniform rod AB has length 3 m and weight W newtons.

The rod is suspended by two light vertical ropes.

One rope is attached to the rod at A and the other rope is attached to the rod at C, where $AC = 2.1 \,\text{m}$.

The rod is in equilibrium in a horizontal position, as shown in Figure 2.

The tension in the rope at C is 350 N.

(a) Show that W = 490

(3)

A particle P of weight 210 N is attached to the rod at a distance d metres from A.

The tension in the rope at C is now 600 N.

The rod remains in equilibrium in a horizontal position.

(b) Find the value of d.

(3)

Particle *P* is removed from the rod.

A particle Q of weight X newtons is now attached at B.

The rod remains in equilibrium in a horizontal position and is now on the point of tilting.

(c) Find the value of X.

(4)



Question 4 continued



Question 4 continued

Question 4 continued	
	Total for Organism Alic 10 - 1
	Total for Question 4 is 10 marks)



5. [*In this question* **i** *and* **j** *are horizontal perpendicular unit vectors and position vectors are given relative to a fixed origin.*]

In a game, a ball B is rolled across a horizontal surface towards a fixed target. The ball is modelled as a particle moving with constant velocity.

At time t = 1 s, the position vector of B is $(-2\mathbf{i} + 5\mathbf{j})$ m.

At time t = 9 s, the position vector of B is $(10\mathbf{i} - 3\mathbf{j})$ m.

(a) Find the velocity of the ball.

(3)

The position vector of the target is (13i - 5j) m.

(b) Use the model to find the distance of B from the target at time t = 7 s.

(4)

Question 5 continued	
(Total for C	Question 5 is 7 marks)



Figure 3

A particle P of mass m is held in equilibrium on a fixed rough inclined plane by a light inextensible string.

The plane is inclined at an angle α to the horizontal, where $\alpha < 45^{\circ}$ The string is inclined to the plane at angle α , as shown in Figure 3.

The string lies in a vertical plane that contains a line of greatest slope of the inclined plane.

When the tension in the string is $0.75 \, mg$, P is on the point of moving up the plane.

(a) Find an expression for the magnitude of the frictional force acting on P, giving your answer in terms of m, g and α .

(3)

The coefficient of friction between P and the plane is $\frac{1}{2}$

(b) Show that

$$\tan \alpha = \frac{2}{5}$$

(6)

The string breaks.

(c) Determine whether P remains at rest. You must justify your reasoning.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued



Question 6 continued		

Question 6 continued	
	(Total for Question 6 is 12 marks)



7.




Figure 4

One end of a light inextensible string is attached to a particle A of mass 2m.

The other end is attached to a particle B of mass 3m.

The string passes over a small smooth fixed pulley.

The string is taut and both straight parts of the string are vertical.

Both particles are held at a distance 0.5 m above a horizontal surface, as shown in Figure 4.

The system is released from rest and *B* moves downwards.

(a) Find the tension in the string in terms of m and g.

(5)

(b) Find the speed of B at the instant it strikes the surface.

(4)

In the subsequent motion, A does not reach the pulley and B does not rebound after it strikes the surface.

(c) Find the time from the instant when the system is released from rest to the instant when A first reaches a height of 1.06 m above the surface.

(6)

Question 7 continued



Question 7 continued	
	(Total for Question 7 is 15 marks)
	TOTAL FOR PAPER IS 75 MARKS

