



Mark Scheme (Results)

January 2023

Pearson Edexcel International Advanced Level
In Pure Mathematics P3 (WMA13) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2023

Question Paper Log Number P72138A

Publications Code WMA13_01_MS_2301

All the material in this publication is copyright

© Pearson Education Ltd 2023

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for this paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:

'M' marks

These are marks given for a correct method or an attempt at a correct method. In Mechanics they are usually awarded for the application of some mechanical principle to produce an equation.

e.g. resolving in a particular direction, taking moments about a point, applying a suvat equation, applying the conservation of momentum principle etc.

The following criteria are usually applied to the equation.

To earn the M mark, the equation

(i) should have the correct number of terms

(ii) be dimensionally correct i.e. all the terms need to be dimensionally correct

e.g. in a moments equation, every term must be a 'force x distance' term or 'mass x distance', if we allow them to cancel 'g' s.

For a resolution, all terms that need to be resolved (multiplied by sin or cos) must be resolved to earn the M mark.

M marks are sometimes dependent (DM) on previous M marks having been earned.

e.g. when two simultaneous equations have been set up by, for example, resolving in two directions and there is then an M mark for solving the equations to find a particular quantity – this M mark is often dependent on the two previous M marks having been earned.

'A' marks

These are dependent accuracy (or sometimes answer) marks and can only be awarded if the previous M mark has been earned. e.g. M0 A1 is impossible.

'B' marks

These are independent accuracy marks where there is no method (e.g. often given for a comment or for a graph).

A few of the A and B marks may be f.t. – follow through – marks.

3. General Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol \checkmark will be used for correct ft
- cao – correct answer only
- cso – correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC – special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp – decimal places
- sf – significant figures
- * – The answer is printed on the paper
- \square – The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao), unless shown, for example as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks are treated as A ft, but manifestly absurd answers should never be awarded A marks.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles)

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$ leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$ leading to $x = \dots$

2. Formula

Attempt to use the correct formula (with values for a , b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0, q \neq 0$, leading to $x = \dots$

Method mark for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values. Where the formula is not quoted, the method mark can be gained by implication from correct working with values but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Question Number	Scheme	Marks
1(a)	$f(x) \leq 9$	B1
		(1)
(b)	$fg(1.5) = f\left(\frac{3}{2 \times 1.5 + 1}\right) = 9 - \left(\frac{3}{2 \times 1.5 + 1}\right)^2$	M1
	$= \frac{135}{16}$	A1
		(2)
(c)	$g(x) = \frac{3}{2x+1} \Rightarrow g^{-1}(x) = \frac{3-x}{2x}$	M1 A1
	$0 < x \leq 3$	B1
		(3)
		Total 6

(a)

B1: Correct range. Allow equivalent notation e.g. $y \leq 9, f \leq 9, y \in (-\infty, 9]$ but not $x \leq 9$.

Condone just " ≤ 9 " and "less than or equal to 9"

(b)

M1: Full attempt at method to find $fg(1.5)$ condoning slips. Implied by a correct answer or 8.44 e.g. For a correct order of operations so requires an attempt to apply $g(1.5)$ first and then f to their $g(1.5)$

Also allow for an attempt to substitute $x = 1.5$ into $9 - \left(\frac{3}{2x+1}\right)^2$ condoning slips such as substituting

$$x = 1.5 \text{ into } 9 - \frac{3}{(2x+1)^2}$$

(c)

M1: Changes the subject of $y = \frac{3}{2x+1}$ and obtains $x = \frac{3 \pm y}{2y}$ or $x = \frac{3}{2y} \pm \frac{1}{2}$ or equivalent.

Alternatively changes the subject of $x = \frac{3}{2y+1}$ and obtains $y = \frac{3 \pm x}{2x}$ or $y = \frac{3}{2x} \pm \frac{1}{2}$ or equivalent.

A1: $g^{-1}(x) = \frac{3-x}{2x}$, $g^{-1}(x) = \frac{1}{2} \left(\frac{3-x}{x} \right)$, $g^{-1}(x) = \frac{3}{2x} - \frac{1}{2}$, $g^{-1}: x \mapsto \frac{3-x}{2x}$ or $g^{-1}: x \mapsto \frac{3}{2x} - \frac{1}{2}$

Condone $y = \frac{3-x}{2x}$ o.e and even $g^{-1} = \frac{3-x}{2x}$ but NOT $f^{-1} = \frac{3-x}{2x}$ o.e. ISW after a correct answer.

$$\frac{3-x}{2}$$

Don't for this mark allow fractions/fractions that may be misinterpreted such as $g^{-1}(x) = \frac{x}{2}$ but answers

such as $g^{-1}(x) = \frac{\frac{3}{2} - 1}{2}$ and $y = \frac{\left(\frac{3-x}{x}\right)}{2}$ are clear and unambiguous and can score both marks.

B1: Correct domain. Allow equivalent notation e.g. $x \in (0, 3]$ but not just $(0, 3]$

Question Number	Scheme	Marks
2(a)	$R = \sqrt{5}$	B1
	$\tan \alpha = \frac{2}{1} \Rightarrow \alpha = \dots$	M1
	$\alpha = 1.107$	A1
		(3)
(b)(i)	$\text{Max} = 3 + 7\sqrt{5}$	B1ft
(b)(ii)	$(2x - "1.107") = \pi \Rightarrow x = \dots$	M1
	$\Rightarrow x = \frac{\pi + "1.107"}{2} = 2.12$	A1
		(3)
		Total 6

(a)

B1: Correct exact value (Condone $R = \pm\sqrt{5}$).

isw after a correct answer. e.g. $R = \sqrt{5} = 2.24$

M1: Allow for: $\tan \alpha = \pm \frac{2}{1}$, $\tan \alpha = \pm \frac{1}{2}$, $\cos \alpha = \pm \frac{1}{"R"}$, $\sin \alpha = \pm \frac{2}{"R"}$ leading to a value for α

If no method is shown imply by the sight of awrt 1.1 rads or awrt 63°

A1: awrt 1.107

(b)(i)

B1ft: Award for $3 + 7 \times \text{their } R$ where $R > 0$.

Also follow through on decimal answers from (a) e.g.

Condone solutions such as $3 - 7 \times -\sqrt{5} = 18.65\dots$

(b)(ii)

M1: For an attempt to solve $(2x \pm "1.107") = \pi$ or $(2x \pm "1.107") = -\pi$. May be implied by awrt 2.12

Condone a bracketing slip $\cos(2(x \pm "1.107")) = -1 \Rightarrow 2(x \pm "1.107") = \pi \Rightarrow x = \dots$ (condone $-\pi$ as above)

Also condone an attempt to solve $(2x \pm "63^\circ") = 180^\circ$ but not mixed units, e.g. $(2x \pm "1.107") = 180^\circ$

A1: For awrt 2.12. This cannot be given in a list, the 2.12 must be selected.

Question Number	Scheme	Marks
3(a)	$\log_{10} y = \frac{5}{16}x + 1.5$	M1A1
		(2)
(b)	$\log_{10} y = \frac{5}{16}x + 1.5 \Rightarrow y = 10^{\frac{5}{16}x + 1.5}$	M1
	$\Rightarrow y = 10^{\frac{5}{16}x} \times 10^{1.5}$	M1
	$y = 31.6 \times 2.05^x$	A1
		(3)
		Total 5

(a)

M1: Scored for a complete attempt to get the equation of the line condoning $\log_{10} y \leftrightarrow y \leftrightarrow l$ and an incorrect sign on the gradient. So, allow for $(\log_{10} y) = \pm \frac{1.5}{4.8}x + 1.5$ o.e. and for $\frac{y-0}{x+4.8} = \pm \frac{1.5}{4.8}$

If this is attempted via simultaneous equations the mark is scored when the candidate reaches $m = \pm 0.3125$
 $c = 1.5$

A1: Correct equation. e.g. $\log_{10} y = \frac{5}{16}x + 1.5$ or equivalent such as $16\log_{10} y = 5x + 24$

The \log_{10} must not appear as "ln" but allow as "log" or "lg"

(b)

Main Method: Starting with their $\log_{10} y = mx + c$

M1: "Removes" the logs in their equation. e.g. $\log_{10} y = "m"x + "c" \Rightarrow y = 10^{m"x + "c"}$

M1: "Correct" strategy to obtain values of k and b or else proceeding correctly to a form $y = 10^{\frac{5}{16}x} \times 10^{1.5}$

Allow for $k = 10^{1.5}$ ($= 31.6$) and $b = 10^{\frac{5}{16}} = (2.05)$. Note that you may see $10^{1.5}$ as $10\sqrt{10}$

A1: Correct equation produced, $y = 31.6 \times 2.05^x$, and no errors seen.

Condone correct working followed by $k = 31.6, b = 2.05$ with the equation being implied

csothe values must be 31.6 and 2.05, not values rounding to these numbers or exact values like $10\sqrt{10}$.

Note: A solution may be fudged from working similar to the following

$$\log_{10} y = mx + c \Rightarrow y = 10^{mx} + 10^c \Rightarrow y = 31.6 + 2.05^x \Rightarrow y = 31.6 \times 2.05^x$$

This will score Special case: M0 M1 A0

Alternative Method: Starting with $y = kb^x$

M1: Takes logs of both sides and applies at least one correct log law

e.g. $\log_{(10)} kb^x = \log_{(10)} k + \log_{(10)} b^x$, $\log_{(10)} b^x = x \log_{(10)} b$

M1: "Correct" strategy to obtain values for k and b from their $y = mx + c$

So $\log_{10} k = "c" \Rightarrow k = 10^c$ and $\log_{10} b = "m" \Rightarrow b = 10^m$

A1: Correct equation produced $y = 31.6 \times 2.05^x$ and no errors seen.

Condone correct working followed by $k = 31.6, b = 2.05$ with the equation being implied

cs0the values must be 31.6 and 2.05, not values rounding to these numbers.

Correct answers for k and b without any working scores M0 M1 A0. Instructions on the paper state that they should show sufficient working to make their method clear.

Question Number	Scheme	Marks
4(a)	Any correct constant, so for $A = 2$ or $B = 3$ or $C = -1$ or $D = 5$	B1
	$2x^4 + 15x^3 + 35x^2 + 21x - 4 = Ax^2(x+3)^2 + Bx(x+3)^2 + C(x+3)^2 + D$ $\Rightarrow A = \dots, B = \dots, C = \dots, D = \dots$ <p style="text-align: center;">or</p> $2x^4 + 15x^3 + 35x^2 + 21x - 4 \div (x^2 + 6x + 9) = \dots x^2 + \dots x + \dots + \frac{\dots}{(x+3)^2}$	M1
	2 correct of $A = 2, B = 3, C = -1, D = 5$	A1
	$A = 2, B = 3, C = -1, D = 5$	A1
		(4)
(b)	$\int f(x) dx = \int \left(2x^2 + 3x - 1 + \frac{5}{(x+3)^2} \right) dx = \frac{2x^3}{3} + \frac{3x^2}{2} - x - \frac{5}{x+3} (+c)$	M1A1ftA1
		(3)
		Total 7

(a)

B1: One correct constant or one correct term in $Ax^2 + Bx + C + \frac{D}{(x+3)^2}$

M1: Complete method for finding A, B, C and D

For example substitution/comparing coefficients/long division

Via substitution/comparing coefficients the minimum required is an identity of the correct form (con doning slips) followed by values for A, B, C and D .

See scheme but there are other versions including

$$2x^4 + 15x^3 + 35x^2 + 21x - 4 = (Ax^2 + Bx + C)(x+3)^2 + D \Rightarrow A = \dots, B = \dots, C = \dots, D = \dots$$

Via division look for a divisor of $x^2 + 6x + 9$, a quotient that is a quadratic and a remainder that is either linear or a constant term.

It could be attempted by dividing by $(x+3)$ twice.

FYI, the first division gives $2x^3 + 9x^2 + 8x - 3$ with a remainder of 5

A1: 2 correct constants following the award of M1

For division, when the remainder is a linear term, it would be scored for two correct of $Ax^2 + Bx + C$

A1: All correct following the award of M1

(b)

M1: $\int \frac{D}{(x+3)^2} dx \rightarrow \frac{k}{x+3}$ where k is a constant.

This may be awarded following a term of $\int \frac{\alpha x + D}{(x+3)^2} dx$ following division.

Look for $\int \frac{\alpha x + D}{(x+3)^2} dx$ being correctly split and $\rightarrow \int \frac{\alpha x}{(x+3)^2} dx + \int \frac{D}{(x+3)^2} dx \rightarrow \text{something} + \frac{k}{x+3}$

A1ft: $\int \left(Ax^2 + Bx + C + \frac{D}{(x+3)^2} \right) dx = \frac{Ax^3}{3} + \frac{Bx^2}{2} + Cx - \frac{D}{x+3} (+c) \cdot$

Correct integration following through on their non-zero constants.

Allow this to be scored with A , B , C and D as above or with made up values

A1: All correct with or without “+ c ”. Allow $-1x$ for $-x$

Question Number	Scheme	Marks
5(a) Way One	$\cot^2 x - \tan^2 x \equiv \frac{\cos^2 x}{\sin^2 x} - \frac{\sin^2 x}{\cos^2 x} \equiv \frac{\cos^4 x - \sin^4 x}{\sin^2 x \cos^2 x}$	M1
	$\equiv \frac{(\cos^2 x - \sin^2 x)(\cos^2 x + \sin^2 x)}{\sin^2 x \cos^2 x} \equiv \frac{\cos 2x}{\dots} \quad \text{or} \quad \frac{\dots}{\left(\frac{1}{2} \sin 2x\right)^2}$	dM1
	$\equiv \frac{\cos 2x}{\left(\frac{1}{2} \sin 2x\right)^2}$	A1
	$\equiv 4 \frac{\cos 2x}{\sin 2x \sin 2x} \equiv 4 \cot 2x \operatorname{cosec} 2x^*$	A1*
	(4)	
(b)	$4 \cot 2\theta \operatorname{cosec} 2\theta = 2 \tan^2 \theta \Rightarrow \cot^2 \theta - \tan^2 \theta = 2 \tan^2 \theta \Rightarrow \cot^2 \theta - 3 \tan^2 \theta = 0$	M1
	$\cot^2 \theta - 3 \tan^2 \theta = 0 \Rightarrow \frac{1}{\tan^2 \theta} - 3 \tan^2 \theta = 0 \Rightarrow \tan^4 \theta = \frac{1}{3}$	A1
	$\tan^4 \theta = \frac{1}{3} \Rightarrow \tan \theta = \pm \sqrt[4]{\frac{1}{3}} = \pm 0.7598.. \Rightarrow \theta = \dots$	M1
	$\theta = \arctan 0.65, -0.65$	A1A1
	(5)	
	Total 9	

(a) Way One LHS to RHS

M1: Changes the LHS to $\sin x$ and $\cos x$ and attempts to make a single fraction using a correct common denominator. Condone errors/slips on the numerator

dM1: Attempts/ applies

- Either $\cos^4 x - \sin^4 x = (\cos^2 x - \sin^2 x)(\cos^2 x + \sin^2 x) = \cos^2 x - \sin^2 x = \cos 2x$ on the numerator
- Or $\sin 2x = 2 \sin x \cos x$ to the denominator condoning bracketing slip

A1: Applies both of the above correctly to achieve a correct expression in terms of $\cos 2x$ and $\sin 2x$

A1*: Reaches the right hand side with sufficient working shown. Expect to see $\sin^2 2x$ split into $\sin 2x \sin 2x$
Penalise consistent (not once or twice) use of poor notation on this mark only.

Examples $\cos^2 x \leftrightarrow \cos x^2$, $\sin^2 \leftrightarrow \sin^2 x$, constantly switching between $x \leftrightarrow \theta$

(b) Allow use of $x \leftrightarrow \theta$

M1: Uses part (a) and attempt to collect terms. (See Appendix III for ways not using part (a))

See below for equations in $\sin \theta$ or $\cos \theta$ where this mark is awarded for an equation in just $\sin \theta$ or $\cos \theta$

A1: Reaches a correct equation in a single term, usually $\tan \theta$. Look for $\tan^4 \theta = \frac{1}{3}$ o.e. such as $3 \tan^4 \theta = 1$

Other correct intermediate forms are $2 \sin^4 \theta + 2 \sin^2 \theta - 1 = 0$ and $2 \cos^4 \theta - 6 \cos^2 \theta + 3 = 0$

M1: Takes the 4th root of their $\frac{1}{3}$ (o.e) and uses \tan^{-1} (you may need to check) to obtain at least one value for θ

For the other intermediate forms look for working such as $\sin^2 \theta = \frac{\sqrt{3}-1}{2} \Rightarrow \sin \theta = \sqrt{\frac{\sqrt{3}-1}{2}} \Rightarrow \theta = \dots$

A1: Either awrt 0.65 or awrt -0.65. Allow either answer in degrees, so awrt $\pm 37.2^\circ$

A1: Both answers in radians, awrt ± 0.65 , and no extras in range

There are many different ways to do part (a). Generally, this is how they will be marked.

Most cases can be aligned to one of the three cases.

RHS to LHS

(a) Way 2	$4 \cot 2x \operatorname{cosec} 2x \equiv 4 \frac{\cos 2x}{\sin 2x} \times \frac{1}{\sin 2x} \equiv \frac{4(\cos^2 x - \sin^2 x)}{\dots} \text{ or } \frac{\dots}{4 \sin^2 x \cos^2 x}$	M1
	$\frac{\cos^2 x - \sin^2 x}{\sin^2 x \cos^2 x} \equiv \frac{1}{\sin^2 x} - \frac{1}{\cos^2 x} \equiv \operatorname{cosec}^2 x - \sec^2 x$	dM1A1
	$\equiv 1 + \cot^2 x - 1 - \tan^2 x \equiv \cot^2 x - \tan^2 x^*$	A1*

M1: Changes to sin 2x and cos 2x or tan 2x and sin 2x and attempts single angles in sin x and cos x

dM1: Changes to single angles throughout and splits into 2 separate fractions (which don't need to be simplified)

A1: Correct expression in terms of the single angles cosec x and sec x

A1*: Reaches the left hand side with sufficient working shown

Working on both sides: One possible way

(a) Way 3	$\cot^2 x - \tan^2 x \equiv 4 \cot 2x \operatorname{cosec} 2x$ $\frac{\cos^2 x}{\sin^2 x} - \frac{\sin^2 x}{\cos^2 x} \equiv 4 \times \frac{\cos 2x}{\sin 2x} \times \frac{1}{\sin 2x}$ $\cos^4 x - \sin^4 x \equiv 4 \sin^2 x \cos^2 x \frac{\cos 2x}{\sin 2x} \times \frac{1}{\sin 2x}$	M1
	$\cos^4 x - \sin^4 x \equiv 4 \times \sin^2 x \cos^2 x \frac{(\cos^2 x - \sin^2 x)}{(2 \sin x \cos x)^2}$	dM1A1
	$(\cos^4 x - \sin^4 x) \equiv (\cos^2 x - \sin^2 x)$ $(\cos^2 x - \sin^2 x) \frac{(\cancel{\cos^2 x + \sin^2 x})}{1} \equiv (\cos^2 x - \sin^2 x) \text{ Hence true}$	A1*

M1: Changes to sin x, cos x, sin 2x and cos 2x and attempts to cross multiply

dM1: Applies

- either $\cos^2 x - \sin^2 x = \cos 2x$ to the numerator
- or $\sin 2x = 2 \sin x \cos x$ to the denominator

A1: Correct identity in terms of cos x and sin x

A1*: Reaches a point where both sides are equal and makes a minimal comment

Example of how you mark a "different" approach.

Question Number	Scheme	Marks
6(a)	(i) $P(0, 3a)$	B1
	(ii) $Q(a, 0) \quad R\left(\frac{7}{3}a, 0\right)$	B1 B1
	(iii) $S\left(\frac{5}{3}a, -2a\right)$	B1
		(4)
(b)	$3x - 5a - 2a = x - 2a \Rightarrow x = \dots$ or $-(3x - 5a) - 2a = -(x - 2a) \Rightarrow x = \dots$	M1
	$x = \frac{5}{2}a \quad \textbf{or} \quad x = \frac{1}{2}a$	A1
	$3x - 5a - 2a = x - 2a \Rightarrow x = \dots$ and $-(3x - 5a) - 2a = -(x - 2a) \Rightarrow x = \dots$	dM1
	$x = \frac{5}{2}a \quad \textbf{and} \quad x = \frac{1}{2}a$	A1
		(4)
		Total 8

(a) Question states simplest form so don't accept e.g. $\frac{10}{6}a$ for $\frac{5}{3}a$

(i)

B1: For $(0, 3a)$. Condone just $y = 3a$ as long as there isn't a value for x coordinate (apart from 0). Note that either of $P = 3a$ or $(3a, 0)$ is B0 unless $y = 3a$ is previously seen

(ii)

B1: For either coordinate $(a, 0)$ **or** $\left(\frac{7}{3}a, 0\right)$ which may or may not be linked correctly to Q and R

Condone just $x = a$ ($x = 1a$) or $x = \frac{7}{3}a$ as long as there isn't a y coordinate (that isn't 0).

Note that an answer such as $Q = a$ or $(0, a)$ is B0 unless $x = a$ is previously seen.

Likewise an answer such as $R = \frac{7}{3}a$ or $(0, 7a/3)$ is B0 unless $x = 7a/3$ is previously seen

B1: For $(a, 0)$ **and** $\left(\frac{7}{3}a, 0\right)$. Allow the coordinates to be given separately. They do not need to be linked to the

correct letter. Condone just $x = a$ ($x = 1a$) and $x = \frac{7}{3}a$ as long as there isn't a y coordinate (that isn't 0).

SC: $Q = a$ **AND** $R = \frac{7}{3}a$ or vice versa is B1 B0

(iii)

B1: $\left(\frac{5}{3}a, -2a\right)$. Allow this to be given as $x = \dots, y = \dots$

(b)

M1: Attempts to solve one **correct** equation. It must be an equation without moduli and be found correctly
For example, look for

- Either taking both positive aspects of the modulus equation $|3x - 5a| - 2a = |x - 2a|$
So $3x - 5a - 2a = x - 2a$ or exact equivalent such as $3x - 7a = x - 2a \Rightarrow x = \dots$
Condone for this mark $\Rightarrow a = \dots$
- Or taking both negative aspects of the modulus equation $|3x - 5a| - 2a = |x - 2a|$
So $-3x + 5a - 2a = -x + 2a$ or exact equivalent such as $-3x + 3a = -x + 2a \Rightarrow x = \dots$
Condone for this mark $\Rightarrow a = \dots$
The $2a$ may be moved over first so it is possible to solve $|3x - 5a| = |x - 2a| + 2a$
So both positive will give an equation $3x - 5a = x - 2a + 2a$ or $3x - 5a = x$
or both negative would give $-3x + 5a = -x + 2a + 2a$ or $-3x + 5a = -x + 4a$

A1: One correct value of x which must be found from a correctly produced equation

dM1: Attempts to solve both of these equations.

There may be other equations which should be ignored for this mark

A1: Both correct values of x (coming from correct equations) with no other values given.

Also allow if 4 answers are found, for example, followed by the 2 correct answers being chosen and the other 2 incorrect answers discarded.

Question Number	Scheme	Marks
7(a)	$x = 3 \tan\left(y - \frac{\pi}{6}\right) \Rightarrow \frac{dx}{dy} = 3 \sec^2\left(y - \frac{\pi}{6}\right)$	B1
	$\Rightarrow \frac{dy}{dx} = \frac{1}{3 \sec^2\left(y - \frac{\pi}{6}\right)}$	M1
	$\frac{1}{3 \left(1 + \tan^2\left(y - \frac{\pi}{6}\right)\right)} = \frac{1}{3 \left(1 + \left(\frac{x}{3}\right)^2\right)}$	dM1
	$= \frac{3}{x^2 + 9}$	A1
		(4)
(b)	$y = \frac{\pi}{3} \Rightarrow x = 3 \tan \frac{\pi}{6} = \sqrt{3}$	B1
	$x = \sqrt{3} \Rightarrow \frac{dy}{dx} = \frac{3}{(\sqrt{3})^2 + 9}$ or $y = \frac{\pi}{3} \Rightarrow \frac{dy}{dx} = \frac{1}{3 \sec^2\left(\frac{\pi}{3} - \frac{\pi}{6}\right)} = \dots$	M1
	$y - \frac{\pi}{3} = \frac{1}{4}(x - \sqrt{3})$	dM1
	$y = 0 \Rightarrow 0 - \frac{\pi}{3} = \frac{1}{4}(x - \sqrt{3}) \Rightarrow x = \dots$	ddM1
	$x = \sqrt{3} - \frac{4\pi}{3}$	A1
		(5)
		Total 9

(a)

B1: Correct derivative, including correct lhs. Condone $\frac{dx}{dy} = k \sec^2\left(y - \frac{\pi}{6}\right)$ where k is a constant.

M1: Either (1) attempts to apply the reciprocal rule $\frac{dy}{dx} = 1 \div \frac{dx}{dy}$.

Don't be too concerned by the position of the constant, it is the function and variable that is important.

Or (2) attempts to apply $\sec^2\left(y - \frac{\pi}{6}\right) = 1 + \tan^2\left(y - \frac{\pi}{6}\right)$ with $\tan\left(y - \frac{\pi}{6}\right)$ being replaced by $\frac{x}{3}$ to get $\frac{dx}{dy}$ in terms of x

dM1: Attempts both (1) and (2) to obtain $\frac{dy}{dx}$ in terms of x

A1: Correct answer

(b)

B1: Correct value for x . Condone awrt 1.73. This can be awarded from the sight of this value in an equation

M1: Uses a correct method to find the value of $\frac{dy}{dx}$ which may be a decimal.

Note that this can be scored either from answer to part (a) e.g. $x = \sqrt{3} \Rightarrow \frac{dy}{dx} = \frac{3}{(\sqrt{3})^2 + 9}$

Or via their $\frac{dx}{dy} = 3 \sec^2 \left(\frac{\pi}{3} - \frac{\pi}{6} \right)$ followed by an attempt at the reciprocal

dM1: Correct straight line method for the tangent at $\left(\sqrt{3}, \frac{\pi}{3} \right)$ using their correctly found m .

It is dependent upon having found the gradient and x value using a correct method.

If they use the form $y = mx + c$ they must proceed as far as $c = \dots$

ddM1: Uses $y = 0$ to find x . It is dependent upon having scored the previous mark

A1: Correct value or exact equivalent for example $\frac{3\sqrt{3} - 4\pi}{3}$

Alt (a) via arctan

B1: $x = 3 \tan \left(y - \frac{\pi}{6} \right) \Rightarrow y = \frac{\pi}{6} + \arctan \left(\frac{x}{3} \right) \Rightarrow \frac{dy}{dx} = 0 + \frac{1}{1 + \left(\frac{x}{3} \right)^2} \times \dots$ where ... could be any values even 1

M1: $\frac{dy}{dx} = 0 + \frac{1}{1 + \left(\frac{x}{3} \right)^2} \times \dots$ where ... could be any value even 1

dM1: $\frac{dy}{dx} = 0 + \frac{1}{1 + \left(\frac{x}{3} \right)^2} \times \frac{1}{3}$

A1*: $\frac{dy}{dx} = \frac{3}{x^2 + 9}$

Alt (a) via compound angle identity they could pick up the first two marks

$$x = \frac{3 \left(\tan y - \tan \frac{\pi}{6} \right)}{1 + \tan \frac{\pi}{6} \tan y} \Rightarrow \frac{dx}{dy} = \frac{\left(1 + \tan \frac{\pi}{6} \tan y \right) \times 3 \sec^2 y - 3 \left(\tan y - \tan \frac{\pi}{6} \right) \times \tan \frac{\pi}{6} \sec^2 y}{\left(1 + \tan \frac{\pi}{6} \tan y \right)^2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\left(1 + \tan \frac{\pi}{6} \tan y \right)^2}{\left(1 + \tan \frac{\pi}{6} \tan y \right) \times 3 \sec^2 y - 3 \left(\tan y - \tan \frac{\pi}{6} \right) \times \tan \frac{\pi}{6} \sec^2 y}$$

Question Number	Scheme	Notes
8	$\int (2 \cos x - \sin x)^2 dx = \int (4 \cos^2 x - 4 \sin x \cos x + \sin^2 x) dx$	M1
	$\int 4 \sin x \cos x dx = \int 2 \sin 2x dx = -\cos 2x$ or $\int 4 \sin x \cos x dx = -2 \cos^2 x \quad \text{or} \quad 2 \sin^2 x$	M1
	$\int (4 \cos^2 x + \sin^2 x) dx = \int (1 + 3 \cos^2 x) dx = \int \left(1 + 3 \left(\frac{\cos 2x + 1}{2} \right) \right) dx$ or $\int (4 \cos^2 x + \sin^2 x) dx = \int \left(4 \left(\frac{\cos 2x + 1}{2} \right) + \frac{1 - \cos 2x}{2} \right) dx$	M1
	$\int (2 \cos x - \sin x)^2 dx = \frac{3}{4} \sin 2x + \cos 2x + \frac{5}{2} x (+c)$ or $\int (2 \cos x - \sin x)^2 dx = \frac{3}{4} \sin 2x + 2 \cos^2 x + \frac{5}{2} x (+c)$ or $\int (2 \cos x - \sin x)^2 dx = \frac{3}{4} \sin 2x - 2 \sin^2 x + \frac{5}{2} x (+c)$	A1A1
		(5)
		Total 5

Condone changes in variables throughout this solution as long as the answer is given in terms of x

M1: Expands to the form $p \cos^2 x + q \sin x \cos x + r \sin^2 x$

M1: Correct strategy for **integrating** $q \sin x \cos x$ (i.e. obtains $k \cos 2x$ or $k \sin^2 x$ or $k \cos^2 x$)

M1: Correct strategy for rewriting $p \cos^2 x + r \sin^2 x$ **into a form that can be integrated.**

Score for one of

- writes in terms of just $\sin^2 x$ and then uses $\sin^2 x = \frac{\pm 1 \pm \cos 2x}{2}$
- writes in terms of just $\cos^2 x$ and then uses $\cos^2 x = \frac{\pm 1 \pm \cos 2x}{2}$
- writes both $\sin^2 x$ and $\cos^2 x$ in terms of $\cos 2x$ with **at least one** of these via use of a correct/allowable form. That is $\sin^2 x = \frac{\pm 1 \pm \cos 2x}{2}$ or $\cos^2 x = \frac{\pm 1 \pm \cos 2x}{2}$

A1: Integrates and achieves 2 correct terms (of the 3 required terms)

NB: An unsimplified expression is acceptable for this mark so please check carefully. e.g. $2x + \dots + \frac{1}{2}x$ counts as one correct term.

A1: Correct simplified integration (+ c not required).

An alternative solution is via " $R \cos$ " or " $R \sin$ " but the last two marks are unlikely to be achieved due to the fact that an exact answer is difficult to arrive at.

Question Number	Scheme	Notes
8	$\int (2 \cos x - \sin x)^2 dx = \int 5 \cos^2 (x + 0.464) dx$	M1
	$= \int \frac{5 \{ \cos (2x + 0.927) + 1 \}}{2} dx$	M1
	$\frac{5 \sin (2x + 0.927)}{4} + \frac{5}{2} x$	M1
	$\frac{3 \sin 2x}{4} + \cos 2x + \frac{5}{2} x + c$	A1A1
		(5)
		Total 5

M1: Writes $2 \cos x - \sin x$ in the form $R \cos(x \pm \alpha)$ o.e. and squares.

Requires a full method so for the form $R \cos(x \pm \alpha)$

- requires $R^2 = 2^2 + 1^2$
- requires $\tan \alpha = \pm \frac{1}{2} \Rightarrow \alpha = \dots$ in radians but condone in degrees

M1: Correct strategy for writing $\cos^2(x \pm \alpha)$ into a form that can be integrated using the double angle formula

$$\cos^2(x \pm \alpha) \rightarrow \frac{\pm 1 \pm \cos(2x \pm 2\alpha)}{2} \text{ but condone } \cos^2(x \pm \alpha) \rightarrow \frac{\pm 1 \pm \cos(2x \pm \alpha)}{2}$$

α should be in radians but condone in degrees

M1: Correct strategy for writing $\cos^2(x \pm \alpha)$ into a form that can be integrated using the double angle formula

$$\cos^2(x \pm \alpha) \rightarrow \frac{\pm 1 \pm \cos(2x \pm 2\alpha)}{2} \text{ but condone } \cos^2(x \pm \alpha) \rightarrow \frac{\pm 1 \pm \cos(2x \pm \alpha)}{2}$$

α should be in radians but condone in degrees

M1: It is for integrating $\cos(2x \pm \delta) \rightarrow \pm \sin(2x \pm \delta)$ following use of an acceptable double angle formula

A1: 2 correct terms of $\frac{3 \sin 2x}{4} + \cos 2x + \frac{5}{2} x + c$

A1: Fully correct and simplified (+ c not required) or $\frac{5 \sin(2x + 2 \arctan 0.5)}{4} + \frac{5}{2} x$

Question Number	Scheme	Notes
8	$\int (2 \cos x - \sin x)^2 \, dx = \int 5 \sin^2 (x - 1.107) \, dx$	M1
	$= \int \frac{5 \{1 - \cos(2x - 2.21)\}}{2} \, dx$	M1
	$= \frac{5}{2} x - \frac{5 \sin(2x - 2.21)}{4}$	M1
	$= \frac{5}{2} x + \frac{3 \sin 2x}{4} + \cos 2x + c$	A1A1
		(5)
		Total 5

Question Number	Scheme	Marks
9(a)	$y = \sqrt{3+4e^{x^2}} = (3+4e^{x^2})^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \frac{1}{2}(3+4e^{x^2})^{-\frac{1}{2}} \times 8xe^{x^2}$	M1
	$= 4xe^{x^2} (3+4e^{x^2})^{-\frac{1}{2}}$	A1
		(2)
(b)	$\frac{(3+4e^{x^2})^{\frac{1}{2}}}{x} = 4xe^{x^2} (3+4e^{x^2})^{-\frac{1}{2}}$	M1
	$\frac{(3+4e^{x^2})}{x} = 4xe^{x^2}$	dM1
	$4x^2e^{x^2} - 4e^{x^2} - 3 = 0^*$	A1*
		(3)
(c)	$f(x) = 4x^2e^{x^2} - 4e^{x^2} - 3 \Rightarrow f(1) = -3 \text{ AND } f(2) = 652 \dots$	M1
	Change of sign and $f(x)$ is continuous hence root in (1, 2)	A1
		(2)
(d)	$4x^2e^{x^2} - 4e^{x^2} - 3 = 0 \Rightarrow x^2 = \frac{4e^{x^2} + 3}{4e^{x^2}} = \frac{4+3e^{-x^2}}{4} \Rightarrow x = \frac{1}{2}\sqrt{4+3e^{-x^2}}^*$	B1*
		(1)
(e)(i)	$x_1 = 1 \Rightarrow x_2 = \frac{1}{2}\sqrt{4+3e^{-1}}$	M1
	$x_3 = 1.0997$	A1
(ii)	$\alpha = 1.1051$	A1
		(3)
		Total 11

(a)

M1: Differentiates using the chain rule to obtain $kxe^{x^2} (3+4e^{x^2})^{-\frac{1}{2}}$ OR $ke^{x^2} (3+4e^{x^2})^{-\frac{1}{2}}$

If they are using implicit differentiation accept $\frac{kxe^{x^2}}{y}$ or $\frac{ke^{x^2}}{y}$

A1: Correct derivative in simplest form. Also award for $\frac{dy}{dx} = \frac{4xe^{x^2}}{y}$ o.e. simplified answer

(b)

M1: Sets their $\frac{dy}{dx} = \frac{(3+4e^{x^2})^{\frac{1}{2}}}{x}$ or equivalent such as use of $y = mx$ with $m =$ their $\frac{dy}{dx}$ and $y = (3+4e^{x^2})^{\frac{1}{2}}$.

Allow with another variable $x \rightarrow \alpha$

dM1: Multiplies up to eliminate the square root. Allow a consistent use of another variable $x \rightarrow \alpha$

Dependent upon the previous M1 and a suitable $\frac{dy}{dx}$, that is one of the form that scored an M1 in part (a)

A1*: Correct proof

An alternative method may be seen using the equation of the line with gradient $\frac{dy}{dx}$ and point $\left(\alpha, \sqrt{3+4e^{\alpha^2}}\right)$

M1: For $y - \sqrt{3+4e^{\alpha^2}} = \left. \frac{dy}{dx} \right|_{x=\alpha} (x - \alpha)$ and sets $(x, y) = (0, 0)$

dM1: Proceeds as the main method with the same condition on the gradient

A1*: Correct proof. Condone with variable α

(c)

M1: Attempts both $f(1)$ and $f(2)$ (or tighter) and obtains at least one value correct to 1sf rounded or truncated.

Condone $f(2) = 12e^4 - 3$ If f or y is not stated assume that $f(x) = 4x^2e^{x^2} - 4e^{x^2} - 3$

A1: Requires

- both values either correct or rounded/ truncated with accuracy to at least 1 sf
- a reference to a sign change e.g. $f(1) = -3 < 0$, $f(2) = 652 > 0$ or $f(1) \times f(2) < 0$
- a mention of continuity
- a minimal conclusion, e.g. hence root

(d)

B1*: Correct proof.

Look for the following evidence

- the line $4x^2e^{x^2} - 4e^{x^2} - 3 = 0$ but may be implied by $4x^2e^{x^2} = 4e^{x^2} + 3$ o.e.

- the line $x^2 = \frac{4e^{x^2} + 3}{4e^{x^2}}$ o.e. $x^2 = 1 + \frac{3}{4e^{x^2}}$ or $4x^2 = \frac{4e^{x^2} + 3}{e^{x^2}}$ o.e. $4x^2 = 4 + \frac{3}{e^{x^2}}$

- correct square root work leading to the given answer

(e)

M1: Attempts to use the iteration formula.

This may be implied by awrt 1.13 for x_2 or awrt 1.10 for x_3 or sight of a correctly embedded value.

It cannot be awarded for just the value of α

A1: awrt 1.0997

A1: 1.1051 cao following the award M1

Question Number	Scheme	Marks
10(a)	$(F =) 35$	B1
		(1)
(b)	$200 = \frac{350e^{15k}}{9 + e^{15k}} \Rightarrow 1800 + 200e^{15k} = 350e^{15k} \Rightarrow 150e^{15k} = 1800$	M1
	$e^{15k} = \frac{1800}{150} \Rightarrow 15k = \ln 12 \Rightarrow k = \frac{1}{15} \ln 12^*$	dM1A1*
		(3)
(c)	$F = \frac{350e^{kt}}{9 + e^{kt}} \Rightarrow \frac{dF}{dt} = \frac{350ke^{kt}(9 + e^{kt}) - 350e^{kt}(ke^{kt})}{(9 + e^{kt})^2}$	M1A1
	$\frac{3150ke^{kt}}{(9 + e^{kt})^2} = 10 \Rightarrow 315ke^{kt} = 81 + 18e^{kt} + e^{2kt} \Rightarrow e^{2kt} + (18 - 315k)e^{kt} + 81 = 0$	M1
	$e^{2kt} + (18 - 315k)e^{kt} + 81 = 0 \Rightarrow e^{kt} = \frac{315k - 18 \pm \sqrt{(18 - 315k)^2 - 4 \times 81}}{2} \Rightarrow kt = \dots$	M1
	$T = \text{awrt } 5.7, 20.8$	A1
		(5)
		Total 9

(a)

B1: Correct value, 35

(b)

M1: Uses $F = 200$ and $t = 15$ and reaches $Ae^{15k} = B$ where $A \times B > 0$

dM1: Proceeds using correct order of operations to obtain a value for k

Look for $Ae^{15k} = B \Rightarrow e^{15k} = \frac{B}{A} \Rightarrow 15k = \ln \frac{B}{A} \Rightarrow k = \dots$

or Alternatively $Ae^{15k} = B \Rightarrow \ln A + 15k = \ln B \Rightarrow 15k = \ln B - \ln A \Rightarrow k = \dots$

A1*: Correct proof with all necessary steps shown.

Minimum acceptable solution $200 = \frac{350e^{15k}}{9 + e^{15k}} \Rightarrow 150e^{15k} = 1800 \Rightarrow e^{15k} = 12 \Rightarrow k = \frac{1}{15} \ln 12$

(c) Allow the whole of part (c) to be done with the letter k , the exact value for k or using $k = \text{awrt } 0.166$

M1: Correct attempt of the quotient (product or chain) rule.

For the Quotient Rule look for $\frac{dF}{dt} = \frac{Pe^{kt}(9 + e^{kt}) - Qe^{kt}(e^{kt})}{(9 + e^{kt})^2} \quad P, Q > 0$

For the Product Rule look for $\frac{dF}{dt} = Pe^{kt}(9 + e^{kt})^{-1} \pm Qe^{kt}e^{kt}(9 + e^{kt})^{-2} \quad P, Q > 0$

For Chain Rule look for $F = A \pm \frac{B}{9 + e^{kt}} \Rightarrow \frac{dF}{dt} = Qe^{kt}(9 + e^{kt})^{-2} \quad Q \text{ is a constant}$

A1: Correct differentiation, which may be unsimplified.

Allow for an expression in k , with exact $k = \frac{1}{15} \ln 12$, or using $k = \text{awrt } 0.166$

M1: Sets their derivative = 10 and obtains a 3TQ in e^{kt}

It is dependent upon a reasonable attempt to differentiate.

In almost all cases the M1 will have been awarded but condone an attempt following $\frac{dF}{dt} = \frac{vu' + uv'}{v^2}$

M1: Scored for

- solving a 3TQ in e^{kt} by any method including a calculator (you may need to check with accuracy to 2sf rounded or truncated). Note that the equation may be a quadratic in $12^{\frac{1}{15}t}$ instead of $e^{\left(\frac{1}{15} \ln 12\right)t}$
- then taking \ln 's to obtain at least one value for kt

FYI the correct quadratics are;

$$e^{\left(\frac{2}{15} \ln 12\right)t} + (18 - 21 \ln 12)e^{\left(\frac{1}{15} \ln 12\right)t} + 81 = 0,$$

$$\text{or } \left(e^{0.166t}\right)^2 - 34.18e^{0.166t} + 81 = 0 \Rightarrow e^{0.166t} = 2.56, 31.62 \Rightarrow 0.166t = 0.94, 3.45$$

A1: awrt 5.7, 20.8

