Please check the examination details bel	ow before ente	ering your candidate info	ormation	
Candidate surname		Other names		
Centre Number Candidate Nu	umber			
Pearson Edexcel International Advanced Level				
Time 1 hour 30 minutes	Paper reference	WMEO	2/01	
Mathematics				
International Advanced Su	ubsidiar	y/Advanced L	.evel	
Mechanics M2				
You must have: Mathematical Formulae and Statistica	al Tables (Ye	ellow), calculator	Total Marks	

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \,\mathrm{m\,s^{-2}}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any
 working underneath.

 Turn over





1.	A particle of mass $0.5 \mathrm{kg}$ is moving with velocity $(2\mathbf{i} + 4\mathbf{j}) \mathrm{m} \mathrm{s}^{-1}$ when it receives an impulse of $(-4\mathbf{i} + 6\mathbf{j}) \mathrm{N} \mathrm{s}$.
	(a) Find the speed of the particle immediately after it receives the impulse. (5)
	(b) Find the size of the angle between the direction of motion of the particle immediately before it receives the impulse and the direction of motion of the particle immediately after it receives the impulse.
	(3)

Question 1 continued	blank
	Q1
(Total 8 marks)	



2. A car of mass 600 kg tows a trailer of mass 200 kg up a hill along a straight road that is inclined at angle θ to the horizontal, where $\sin \theta = \frac{1}{20}$. The trailer is attached to the car by

a light inextensible towbar. The resistance to the motion of the car from non-gravitational forces is modelled as a constant force of magnitude $150\,\mathrm{N}$. The resistance to the motion of the trailer from non-gravitational forces is modelled as a constant force of magnitude $300\,\mathrm{N}$.

When the engine of the car is working at a constant rate of PkW the car and the trailer have a constant speed of $15\,\mathrm{m\,s^{-1}}$

(a) Find the value of P.

(5)

Later, at the instant when the car and the trailer are travelling up the hill with a speed of $20 \,\mathrm{m\,s^{-1}}$, the towbar breaks. When the towbar breaks the trailer is at the point X. The trailer continues to travel up the hill before coming to instantaneous rest at the point Y. The resistance to the motion of the trailer from non-gravitational forces is again modelled as a constant force of magnitude $300 \,\mathrm{N}$.

(b) Use the work-energy principle to find the distance XY.

- 1	4	1
•	4	1
•		,



	Leave blank
Question 2 continued	



estion 2 continued	

Question 2 continued	Leave blank
	Q2
(T ₀ 4 ₀ 1 0	
(Total 9 marks)	



3. A particle *P* of mass 0.25 kg is moving on a smooth horizontal surface under the action of a single force, **F** newtons.

At time t seconds $(t \ge 0)$, the velocity $\mathbf{v} \, \mathbf{m} \, \mathbf{s}^{-1}$ of P is given by

$$\mathbf{v} = (6\sin 3t)\mathbf{i} + (1 + 2\cos t)\mathbf{j}$$

(a) Find \mathbf{F} in terms of t.

(3)

At time t = 0, the position vector of P relative to a fixed point O is $(4\mathbf{i} - \sqrt{3}\mathbf{j})$ m.

(b) Find the position vector of P relative to O when P is first moving parallel to the vector \mathbf{i} .

(6)

	Leave blank
Question 3 continued	Olalik



Leave blank

Question 3 continued		

Question 3 continued	Leave
Question 5 continued	
	Q3
(Total 9 marks)	



4. Two small balls, A and B, are moving in opposite directions along the same straight line on smooth horizontal ground. The mass of A is 2m and the mass of B is 3m. The balls collide directly. Immediately before the collision, the speed of A is 2u and the speed of B is a. The coefficient of restitution between A and B is a, where a is a in a in

By modelling the balls as particles,

(a) show that the speed of B immediately after the collision is $\frac{1}{5}u(1+6e)$.

(6)

After the collision with ball A, ball B hits a smooth fixed vertical wall which is perpendicular to the direction of motion of B.

The coefficient of restitution between B and the wall is $\frac{5}{7}$

Ball B rebounds from the wall and there is a second direct collision between A and B.

(b) Find the range of possible values of e.

	-		•	,
1	Ι.	Λ	ı	
	٠.	_	r	

	Lo
estion 4 continued	
	1



Leave blank

Question 4 continued

Question 4 continued	blank
	Q4
(Total 10 marks)	



5.




Figure 1

A smooth solid hemisphere is fixed with its flat surface in contact with rough horizontal ground. The hemisphere has centre O and radius 5a.

A uniform rod AB, of length 16a and weight W, rests in equilibrium on the hemisphere with end A on the ground. The rod rests on the hemisphere at the point C, where AC = 12a and angle CAO = a, as shown in Figure 1.

Points A, C, B and O all lie in the same vertical plane.

(a) Explain why
$$AO = 13a$$

(1)

The normal reaction on the rod at C has magnitude kW

(b) Show that
$$k = \frac{8}{13}$$

(3)

The resultant force acting on the rod at A has magnitude R and acts upwards at θ° to the horizontal.

- (c) Find
 - (i) an expression for R in terms of W
 - (ii) the value of θ

(8)

	Lo bl
uestion 5 continued	



Leave blank

Question 5 continued		

Question 5 continued	blank
	Q5
(Total 12 marks)	



6. [The centre of mass of a semicircular arc of radius r is $\frac{2r}{\pi}$ from the centre.]




Figure 2

Uniform wire is used to form the framework shown in Figure 2.

In the framework,

- *ABC* is straight and has length 25*a*
- *ADE* is straight and has length 24*a*
- *ABD* is a semicircular arc of radius 7*a*
- EC = 7a
- angle $AEC = 90^{\circ}$
- the points A, B, C, D and E all lie in the same plane

The distance of the centre of mass of the framework from AE is d.

(a) Show that
$$d = \frac{53}{2(7+\pi)}a$$
 (4)

The framework is freely suspended from A and hangs in equilibrium with AC at angle α° to the downward vertical.

(b) Find the value of α .

(7)



nestion 6 continued	ł
	1

Leave blank

Question 6 continued	

	Leave
Question 6 continued	
	Q6
(Total 11 marks)	



7. A particle P is projected from a fixed point O on horizontal ground. The particle is projected with speed u at an angle α above the horizontal. At the instant when the horizontal distance of P from O is x, the vertical distance of P above the ground is y. The motion of P is modelled as that of a particle moving freely under gravity.

(a) Show that
$$y = x \tan \alpha - \frac{gx^2}{2u^2} \left(1 + \tan^2 \alpha\right)$$
 (6)

A small ball is projected from the fixed point O on horizontal ground. The ball is projected with speed $20 \,\mathrm{m\,s^{-1}}$ at angle θ° above the horizontal. A vertical pole AB, of height $2 \,\mathrm{m}$, stands on the ground with $OA = 10 \,\mathrm{m}$, as shown in Figure 3.




Figure 3

The ball is modelled as a particle moving freely under gravity and the pole is modelled as a rod.

The path of the ball lies in the vertical plane containing O, A and B.

Using the model,

(b) find the range of values of θ for which the ball will pass over the pole.

(3)

Given that $\theta = 40$ and that the ball first hits the ground at the point C

(c) find the speed of the ball at the instant it passes over the pole,

(5)

(d) find the distance OC.

(2)

	Leave blank
Question 7 continued	



Leave blank

Question 7 continued

	Leave blank
Question 7 continued	



TOTAL FOR PAPER IS 75 MARKS

END