



Mark Scheme (Results)

January 2020

Pearson Edexcel International A Level
in Statistics S2 (WST02) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2020

Publications Code WST02_01_MS_2001*

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.



- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme		Marks
1 (a)	$P(H = 6) = \frac{e^{-4} 4^6}{6!}$ or $P(H \leq 6) - P(H \leq 5) = 0.8893 - 0.7851$		M1
	$= 0.10419\dots$	$= 0.1042$ awrt 0.104	A1
			(2)
(b)	$J \sim \text{Po}(8)$		B1
	$P(J \leq 7) - P(J \leq 2) = 0.4530 - 0.0138$		M1
	$= 0.4392$	awrt 0.439	A1
(c)	$K \sim N(28, 28)$		M1
	$P(K > 30) \approx P\left(Z > \frac{30.5 - 28}{\sqrt{28}}\right)$		M1M1A1
	$= P(Z > 0.4724\dots)$		
	$= 1 - 0.6808$		
	$= 0.3192$ (calc 0.3183)	awrt 0.319/0.318	A1
			(5)
(d)(i)	The p(robability)/0.97 is not small oe		B1
			(1)
(ii)	$L \sim \text{Po}(3)$		B1
	$P(L \leq 4) = 0.8153$	awrt 0.815	M1A1
			(3)
	Notes		Total 14
	Correct answers imply all marks in each part of this question.		
1(a)	M1	Allow any value for lambda $\frac{e^{-\lambda} \lambda^6}{6!}$ or $P(H \leq 6) - P(H \leq 5)$	
	A1	awrt 0.104	
(b)	B1	Writing Po(8). This may be implied by a correct answer or sight of awrt 0.453 or awrt 0.0138 or awrt 0.0424 or awrt 0.313	
	M1	$P(J \leq 7) - P(J \leq 2)$ oe or (awrt 0.453 – awrt 0.0138)	
	A1	awrt 0.439	
(c)	M1	Using normal approximation with mean = variance = 28 (May be seen in standardisation which takes priority) or writing N(28,28)	
	M1	$\pm \left(\frac{30.5 \text{ or } 30 \text{ or } 29.5 - \text{their mean}}{\text{their sd}} \right)$	
	M1	If they have not given a mean and a variance, they must be correct here.	
	M1	Writing or using a continuity correction 30 ± 0.5	
	A1	Correct standardisation with 30.5 or awrt 0.47	
	A1	awrt 0.319/0.318	
(d)(i)	B1	Probability is not small (too large). Allow mean \neq variance.	
		Do not allow e.g. ‘np too large/np > 10’ on its own.	
		Ignore extraneous non-contradictory comments.	
(ii)	B1	Writing or using Po(3)	
	M1	Writing or using $P(L \leq 4)$ oe	
	A1	awrt 0.815	

Question	Scheme		Marks
2(a)	$E \sim B(6, 0.35)$		B1
(i)	$P(E = 2) = P(E \leq 2) - P(E \leq 1)$ or $\binom{6}{2} 0.35^2 (1 - 0.35)^4$		M1
	$= 0.6471 - 0.3191$		
	$= 0.328$		awrt 0.328
(ii)	$P(E \geq 4) = 1 - P(E \leq 3)$ or $1 - 0.8826$		M1
	$= 0.1174$		awrt 0.117
			(5)
(b)	$H_0 : p = 0.35 \quad H_1 : p > 0.35$		B1
	$L \sim B(50, 0.35) \quad P(L \geq 25) = 1 - P(L \leq 24)$	$P(L \geq 24) = 0.0396$	M1
	$= 1 - 0.9793$	$P(L \geq 23) = 0.071$	
	$= 0.0207$	CR $L \geq 24$	A1
	Reject H_0 or Significant or 25 does lie in the critical region		dM1
	There is evidence to support Kiyoshi's belief oe or that the proportion/number oe of large eggs has increased after adding the supplement		A1cso
			(5)
(c)	Expected profit before supplement = $0.1174 \times 1.20 + (1 - 0.1174) \times 0.60$		M1
	$= (\pounds)0.67044$		
	$P(X \geq 4) = 0.2553$	awrt 0.255	B1
	Expected profit per box after supplement = $0.2553 \times 1.20 + (1 - 0.2553) \times 0.60 - 0.67044$		M1
	$= (\pounds)0.08274$		A1
	OR Expected profit per box after supplement = $0.2553 \times 1.20 + (1 - 0.2553) \times 0.60 - 0.10$		(M1)
	$= (\pounds)0.65318$		(A1)
	Kiyoshi should not continue to add the supplement (as $0.0827 < 0.10$ or $0.653 < 0.67[0]$)		A1cso
			(5)
			Total 15
	Notes		
2(a)	B1	Using or writing $B(6, 0.35)$ in either part	
(i)	M1	Using or writing $P(E \leq 2) - P(E \leq 1)$ oe or writing $\binom{6}{2} 0.35^2 (1 - 0.35)^4$ oe	
	A1	awrt 0.328	
(ii)	M1	Either writing or using $P(E \geq 4)$ or $1 - P(E \leq 3)$ or $P(E = 4) + P(E = 5) + P(E = 6)$ oe	
	A1	awrt 0.117 (Correct answers imply all previous marks in part (a))	
(b)	B1	Both hypotheses correct with p or π	
	M1	Writing or using $L \sim B(50, 0.35)$ and $1 - P(L \leq 24)$	
		or writing $P(L \geq 24) = 0.0396$ or $P(L \geq 23) = 0.071$ leading to a CR.	
	A1	Condone use of normal approx $M \sim N(17.5, \text{awrt } 11.4)$ and $1 - P(M < 24.5)$ for the M1	
	dM1	awrt 0.0207 or $L \geq 24$ allow any letter	
	dM1	dep on previous M being awarded for a correct statement (condone Accept H_1)	
	A1cso	ft their probability or CR Do not allow contradicting non-contextual comments.	
		All previous marks must be awarded. A correct statement in context.	
		Need bold words . NB award M1A1 for a correct contextual statement on its own.	
(c)	Note:	Some candidates may multiply by n or an integer so allow these multiples throughout.	
	M1	"their (ii) $\times 1.20 + (1 - \text{"their(ii)"}) \times 0.60$	
	B1	awrt 0.255	
	M1	$"p" \times 1.20 + (1 - "p") \times 0.60 - \text{"their } 0.67044"$ or $"p" \times 1.20 + (1 - "p") \times 0.60 - 0.10$ oe	
		where $p > 0.1174$ (do not allow $p = 0.45$ for this mark)	
	A1	awrt $(\pounds)0.083$ or awrt $(\pounds)0.65$	
	A1cso	Dep on all previous marks in (c). Correct conclusion with correct supporting figures.	

Question Number	Scheme		Marks
3(a)	$\frac{3}{4}$		B1
			(1)
(b)	$E(T) = \frac{50+2k}{2} [= 25+k]$		B1
	$Var(T) = \frac{(4k)^2}{12} \left[= \frac{4k^2}{3} \right]$		B1
	$E(T^2) = \frac{4k^2}{3} + (25+k)^2$		M1
	$\frac{7k^2}{3} + 625 + 50k = 918.76$		
	$7k^2 + 150k - 881.28 = 0$		dM1
	$k = \frac{-150 \pm \sqrt{150^2 + 4 \times 7 \times 881.28}}{14}$		dM1
	$k = 4.8$ oe only		A1
			(6)
(c)	$P(T < 25) = \frac{1}{4}$		B1
	B(50, 0.25)		
	$P(X \geq 20) = 1 - P(X \leq 19)$		M1
	$= 1 - 0.9861$		
	$= 0.0139$	awrt 0.0139	A1
			(3)
			Total 10
Notes			
(a)	B1	0.75 oe	
(b)	B1	$E(T) = \frac{50+2k}{2} [= 25+k]$ allow equivalent unsimplified expressions	
	B1	$Var(T) = \frac{(4k)^2}{12} \left[= \frac{4k^2}{3} \right]$ allow equivalent unsimplified expressions	
	M1	Using $Var(T) + [E(T)]^2$ oe e.g. " $\frac{4k^2}{3}$ " = $E(T^2) - (25+k)^2$	
	dM1	Dependent on previous M being awarded. Substituting $E(T^2) = 918.76$, multiplying out and combining like terms leading to a $3TQ = 0$	
	dM1	Dependent on previous M being awarded. A correct method for solving their quadratic – use of formula (allow one slip), completing the square, factorising.	
	A1	Must have 4.8 oe on its own as answer (must reject $k = -26.2$ if seen)	
	ALT	4.8 on its own scores 6 out of 6.	
		For first 4 marks in (b)	
		$\int_{25-k}^{25+3k} t^2 \left(\frac{1}{4k}\right) dt = \left[\frac{t^3}{12k} \right]_{25-k}^{25+3k} \rightarrow \frac{(25+3k)^3}{12k} - \frac{(25-k)^3}{12k} = 918.76$	
		B2 for correct integral (ignore limits), M1 for attempt at integration $t^2 \rightarrow t^3$, dM1 for use of limits and = 918.76, then follow main scheme	
(c)	B1	0.25	
	M1	Writing or using $1 - P(X \leq 19)$	
	A1	awrt 0.0139	

Question Number	Scheme		Marks
4(a)	$\int_1^2 \frac{1}{3} dt + \int_2^4 k(4t^2 - t^3) dt = 1$		M1
	$\left[\frac{1}{3}t \right]_1^2 + \left[k \left(\frac{4t^3}{3} - \frac{t^4}{4} \right) \right]_2^4 = 1$		A1
	$\frac{1}{3} + k \left(\frac{64}{3} - \frac{20}{3} \right) = 1$ or $\frac{44}{3}k = \frac{2}{3}$ leading to $k = \frac{1}{22}$		A1 cso
			(3)
(b)			B1(shape)
			dB1
			(labels)
			(2)
(c)	$\frac{df(t)}{dt} = k(8t - 3t^2)$		B1
	$8t - 3t^2 = 0$		M1
	$t = \frac{8}{3}$ only	awrt 2.67	A1
			(3)
(d)	$\int_1^t \frac{1}{3} dx = \left[\frac{x}{3} \right]_1^t$		M1
	$F(2) + \int_2^t \frac{1}{22} (4x^2 - x^3) dx = \frac{1}{3} + \left[\frac{4x^3}{66} - \frac{x^4}{88} \right]_2^t$		M1
	Or $\int \frac{1}{22} (4t^2 - t^3) dt = \frac{2t^3}{33} - \frac{t^4}{88} + C$ and $F(4) = 1$		
	$F(t) = \begin{cases} 0 & t < 1 \\ \frac{1}{3}t - \frac{1}{3} & 1 \leq t < 2 \\ \frac{2t^3}{33} - \frac{t^4}{88} + \frac{1}{33} & 2 \leq t \leq 4 \\ 1 & \text{otherwise} \end{cases}$		A1 A1 A1
			(5)
(e)	$P(T > 3) = 1 - F(3)$		
	$= 1 - \left[\frac{4 \times 3^3}{66} - \frac{3^4}{88} + \frac{1}{33} \right]$		M1
	$= \frac{67}{264}$ or 0.2537...	awrt 0.254	A1
			(2)
			Total 15

		Notes
4(a)	M1	Adding the two integrals together with correct limits and setting = 1 (may be done in stages) Allow $\frac{1}{3}$ instead of first integral
	A1	Correct integration (again allow $\frac{1}{3}$ instead of first integration)
	A1cso	Must have at least one line of working before the given answer and no errors
(b)	B1	Correct shape with correct curvature Horizontal line, then quadratic (increasing then decreasing as t increases) starting above horizontal line and finishing on horizontal axis. The sketch is not continuous. There should be no solid vertical lines.
	dB1	Fully correct with 1, 2 and 4 each labelled at appropriate place on horizontal axis (Ignore vertical labelling). e.g.
		 
		B0B0 (solid vertical line) B1B1 Condone curvature
(c)	B1	For $k(8t - 3t^2)$
	M1	Putting their differential = 0 ignore missing k
	A1	Allow awrt 2.67 only
(d)	M1	For $\int_1^t \frac{1}{3} dx$ with attempt to integrate. Must have correct limits. Or for integration with +C and use of $F(1) = 0$
	M1	For $F(2) + \int_2^t \frac{1}{22}(4x^2 - x^3) dx$ and attempt to integrate or $\int \frac{1}{22}(4t^2 - t^3) dt = \frac{4t^3}{66} - \frac{t^4}{88} + C$ and using $F(4) = 1$ or $F(2) = \frac{1}{3}$ – must attempt to integrate, have + C
	A1	For 2 nd line of cdf oe (allow < instead of \leq and vice versa ditto > and \geq) (allow any letter to be used for this A1 mark)
	A1	For 3 rd line of cdf oe (allow < instead of \leq and vice versa ditto > and \geq) (allow any letter to be used for this A1 mark)
	A1	All correct and in terms of t including $F(t)$. Allow the otherwise to be for any of the parts but there must be only one. (allow < instead of \leq and vice versa ditto > and \geq)
(e)	M1	Attempting to find $1 - F(3)$ with attempt to use 3 rd line of their $F(t)$ or $\int_3^4 k(4t^2 - t^3) dt$
	A1	$\frac{67}{264}$ oe or awrt 0.254

Question Number	Scheme		Marks
5(a)	$X \sim \text{Po}(4)$		M1
	$P(X = 0) = 0.0183$	$P(X \geq 8) = 0.0511$	
	$P(X \leq 1) = 0.0916$	$P(X \geq 9) = 0.0214$	
	CR $X = 0$ oe	$X \geq 9$ oe	A1A1
			(3)
(b)	3.97%		B1
			(1)
(c)	6 is not in the critical region – the data collected are consistent with Chris’s claim		B1ft
			(1)
(d)	$\lambda = \frac{2n}{9}$		B1
	$1 - P(Y = 0) > 0.9$		M1
	$1 - e^{-\frac{2n}{9}} > 0.9$		
	$e^{-\frac{2n}{9}} < 0.1$		
	$n = 10$ and $e^{-\frac{2n}{9}} = 0.1083...$	or $-\frac{2n}{9} < \ln 0.1$	dM1
	$n = 11$	$e^{-\frac{2n}{9}} = 0.08677...$	
	Therefore $n = 11$		A1 cao
			(4)
(e)	$H_0 : \lambda = 10$ $H_1 : \lambda < 10$		B1
	$[W \sim \text{Po}(10) \quad P(W \leq 5) =] 0.0671$ or CR $W \leq 4$		B1
	Do not reject H_0 or insignificant or 5 does not lie in the critical region		M1
	There is no significant evidence that the mean number/rate of whales has decreased.		A1cso
			(4)
			Total 13
	Notes		
(a)	M1 A1 A1	Writing or using Po(4) (may be implied by one correct CR) Either tail $X = 0$ (allow $X \leq 0$) or $[18 \geq] X \geq 9$ (allow $X > 8$) Allow any letters in place of X Both tails $X = 0$ oe , $[18 \geq] X \geq 9$ oe Allow any letters in place of X SC: $P(X = 0)$ and $P(X \geq 9)$ as final answer to score M1A1A0.	
(b)	B1	awrt 3.97% or awrt 0.0397	
(c)	B1ft	Supports this claim and correct reason. Allow a correct f.t. statement and reason based on their CR	
(d)	B1	writing or using $\frac{2n}{9}$	
	M1	May be implied by $P(Y = 0) < 0.1$ (Allow = in place of <)	
	dM1	Dep on previous M mark for solving $e^{-\lambda} < 0.1$. This may be implied by $n = \text{awrt } 10.4$ Allow for a trial of any n value or $-\frac{2n}{9} < \ln 0.1$ (condone $\frac{2n}{9} = 2.5$) (Allow = in place of <)	
(e)	A1	11 cao (Do not allow $n \geq 11$)	
	B1	Both hypotheses with λ or μ (Allow $H_0 : \lambda = 2$ $H_1 : \lambda < 2$)	
	B1	awrt 0.0671 or $W \leq 4$	
	M1	Correct statement – ft their probability or CR Do not allow contradicting non-contextual comments	
	A1cso	Fully correct solution with conclusion in context must have mean/rate with whales. NB award M1A1 for a correct contextual statement on its own provided previous marks scored	

Question Number	Scheme		Marks
6(a)	$E(X^2) = \int_{-1}^1 \frac{1}{8}(x^4 + 2x^3 + x^2) dx + \int_1^{\frac{11}{3}} \frac{1}{4}x^2 dx$		M1
	$= \left[\frac{1}{8} \left(\frac{x^5}{5} + \frac{2x^4}{4} + \frac{x^3}{3} \right) \right]_{-1}^1 + \left[\frac{x^3}{12} \right]_1^{\frac{11}{3}}$		A1
	$= \frac{1684}{405}$		
	$\text{Var}(X) = \frac{1684}{405} - \left(\frac{31}{18} \right)^2$		dM1
	$= \frac{1931}{1620} \text{ or } 1.1919\dots$	awrt 1.19	A1
			(4)
(b)	$P\left(X < -\frac{1}{2}\right) = \int_{-1}^{-0.5} \frac{1}{8}(x^2 + 2x + 1) dx$	or $1 - \int_{-0.5}^{0.5} \frac{1}{8}(x^2 + 2x + 1) dx$ (gets M2)	M1
	$P\left(X > \frac{1}{2}\right) = \frac{2}{3} + \int_{0.5}^1 \frac{1}{8}(x^2 + 2x + 1) dx$		M1
	$P\left(X < -\frac{1}{2}\right) = \left[\frac{x^3}{24} + \frac{x^2}{8} + \frac{x}{8} \right]_{-1}^{-0.5}$ or $P\left(X > \frac{1}{2}\right) = \frac{2}{3} + \left[\frac{x^3}{24} + \frac{x^2}{8} + \frac{x}{8} \right]_{0.5}^1$		A1
	or $1 - \left[\frac{x^3}{24} + \frac{x^2}{8} + \frac{x}{8} \right]_{-0.5}^{0.5}$		
	$= \frac{83}{96} \text{ or } 0.8645\dots$	awrt 0.865	A1
			(4)
			Total 8
	Notes		
	In parts (a) and (b) a correct answer does NOT imply all marks.		
(a)	M1 A1 dM1 A1	For attempt at $\int x^2 f(x) dx$ for both parts of $f(x)$ added and attempt to integrate $x^n \rightarrow x^{n+1}$ Correct algebraic integration (ignore limits). This mark cannot be implied. dep on previous M1 for “an expression for their $E(X^2)$ ” – $[E(X)]^2$ Values must be substituted here awrt 1.19	
(b)	M1 M1 NB A1 A1	Main scheme method $\int_{-1}^{-0.5} \frac{1}{8}(x^2 + 2x + 1) dx$ oe $\frac{2}{3} + \int_{0.5}^1 \frac{1}{8}(x^2 + 2x + 1) dx$ oe $1 - \int_{-0.5}^{0.5} \frac{1}{8}(x^2 + 2x + 1) dx$ gets M2 One correct integration (may be implied by $\frac{1}{192}, \frac{9}{64}$ or $\frac{55}{64}$)	Alternative method using F(x) $\int_{-1}^x \frac{1}{8}(t^2 + 2t + 1) dt$ or $\int \frac{1}{8}(x^2 + 2x + 1) dx$ with + C Use of $F(-0.5) + (1 - F(0.5))$ oe Note: $F(-0.5) = \frac{1}{192}$ and $F(0.5) = \frac{9}{64}$ $F(x) = \frac{1}{8} \left(\frac{x^3}{3} + x^2 + x + \frac{1}{3} \right)$ (from $-1 < x < 1$) $\frac{83}{96}$ or awrt 0.865 must come from correct working and dependent on all previous marks.

