Please check the examination detail	ls below before en	tering your candidate	information
Candidate surname		Other names	
Centre Number Candidat	te Number		
Pearson Edexcel Int	ernation	nal Advan	ced Level
carson Euckeerine			
Time 1 hour 30 minutes	Paper reference	WFM	01/01
Mathematics			00
International Advanced	l Cubcidia	w/Advance	d Lovel
		y/Advanced	u Levei
Further Pure Mathemat	IICS F I		
You must have: Mathematical Formulae and Stati	stical Tables (Y	ellow), calculato	Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- You should show sufficient working to make your methods clear.
  Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## Advice

- Read each guestion carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.









**(5)** 

1.	$\mathbf{M} = \begin{pmatrix} 3x & 7 \\ 4x + 1 & 2 - x \end{pmatrix}$
	Find the range of values of $x$ for which the determinant of the matrix $M$ is positive.

2	
_	

	blank
Question 1 continued	
	·
	_
	_
	-
	-
	-
	-
	-
	-
	-
	-
	_
	_
	-
	-
	-
	-
	-
	-
	-
	_
	Q1
(Total 5 marks)	,
( * : : : * * : : : : : : : : : : : : :	



2. The complex numbers  $z_1$  and  $z_2$  are given by

$$z_1 = 3 + 5i$$
 and  $z_2 = -2 + 6i$ 

(a) Show  $z_1$  and  $z_2$  on a single Argand diagram.

**(2)** 

- (b) Without using your calculator and showing all stages of your working,
  - (i) determine the value of  $|z_1|$

**(1)** 

(ii) express  $\frac{z_1}{z_2}$  in the form a + bi, where a and b are fully simplified fractions.

**(3)** 

(c) Hence determine the value of  $\arg \frac{z_1}{z_2}$ 

Give your answer in radians to 2 decimal places.

**(2)** 



	Leave
	blank
Question 2 continued	



Leave blank

Question 2 continued	

	bla
(Total 8 marks)	Q2



3.	The parabola	C has e	quation	$y^2 =$	18 <i>x</i>
----	--------------	---------	---------	---------	-------------

The point S is the focus of C

(a) Write down the coordinates of S

**(1)** 

Leave blank

The point P, with y > 0, lies on C

The shortest distance from P to the directrix of C is 9 units.

(b) Determine the exact perimeter of the triangle *OPS*, where *O* is the origin.

Give your answer in simplest form.

**(4)** 






	Leave
	blank
Question 3 continued	
	1



Leave blank

Question 3 continued	

Question 3 continued		blank
		Q3
	(Total 5 marks)	



4. The equation

$$x^4 + Ax^3 + Bx^2 + Cx + 225 = 0$$

where A, B and C are real constants, has

- a complex root 4 + 3i
- a repeated positive real root
- (a) Write down the other complex root of this equation.

(1)

(b) Hence determine a quadratic factor of  $x^4 + Ax^3 + Bx^2 + Cx + 225$ 

(2)

(c) Deduce the real root of the equation.

**(2)** 

(d) Hence determine the value of each of the constants A, B and C

(3)


	Leave
	blank
Question 4 continued	



Leave blank

Question 4 continued

Question 4 continued	blank
	Q4
/T-1-10 1 )	
(Total 8 marks)	



$$\mathbf{P} = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$$

The matrix  $\bf P$  represents the transformation U

(a) Give a full description of U as a single geometrical transformation.

**(2)** 

The transformation V, represented by the  $2 \times 2$  matrix Q, is a reflection in the line y = -x

(b) Write down the matrix  $\mathbf{Q}$ 

**(1)** 

The transformation U followed by the transformation V is represented by the matrix  $\mathbf{R}$ 

(c) Determine the matrix **R** 

**(2)** 

The transformation W is represented by the matrix  $3\mathbf{R}$ 

The transformation W maps a triangle T to a triangle T'

The transformation W' maps the triangle T' back to the original triangle T

(d) Determine the matrix that represents W'

**(3)** 



	Leave
	blank
Question 5 continued	



Leave blank

Question 5 continued	l		

Question 5 continued		blan
		Q5
	(Total 8 marks)	



**6.** The quadratic equation

$$Ax^2 + 5x - 12 = 0$$

where A is a constant, has roots  $\alpha$  and  $\beta$ 

- (a) Write down an expression in terms of A for
  - (i)  $\alpha + \beta$
  - (ii)  $\alpha\beta$

**(2)** 

The equation

$$4x^2 - 5x + B = 0$$

where *B* is a constant, has roots  $\alpha - \frac{3}{\beta}$  and  $\beta - \frac{3}{\alpha}$ 

(b) Determine the value of A

**(3)** 

(c) Determine the value of B

**(3)** 



	bl
Jugation 6 continued	01
Question 6 continued	
	_
	—
	—
	—
	_



Leave blank

Question 6 continued		

Question 6 continued		blank
		<b>Q</b> 6
	(Total 8 marks)	



7. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The rectangular hyperbola H has equation xy = 36

The point P(4, 9) lies on H

(a) Show, using calculus, that the normal to H at P has equation

$$4x - 9y + 65 = 0$$

(4)

The normal to H at P crosses H again at the point Q

(b) Determine an equation for the tangent to H at Q, giving your answer in the form y = mx + c where m and c are rational constants.

**(5)** 

	Leave
	blank
Question 7 continued	



Leave blank

Question 7 continued

Question 7 continued	blank
	<b>Q</b> 7
(Takal 0 maulus)	
(Total 9 marks)	



$$f(x) = 2x^{-\frac{2}{3}} + \frac{1}{2}x - \frac{1}{3x - 5} - \frac{5}{2} \qquad x \neq \frac{5}{3}$$

The table below shows values of f(x) for some values of x, with values of f(x) given to 4 decimal places where appropriate.

х	1	2	3	4	5
f(x)	0.5		-0.2885		0.5834

(a) Complete the table giving the values to 4 decimal places.

**(2)** 

The equation f(x) = 0 has exactly one positive root,  $\alpha$ .

Using the values in the completed table and explaining your reasoning,

(b) determine an interval of width one that contains  $\alpha$ .

**(2)** 

(c) Hence use interval bisection twice to obtain an interval of width 0.25 that contains  $\alpha$ .

**(3)** 

Given also that the equation f(x) = 0 has a negative root,  $\beta$ , in the interval [-1, -0.5]

(d) use linear interpolation once on this interval to find an approximation for  $\beta$ .

Give your answer to 3 significant figures.

**(3)** 



uestion 8 continued	



Leave blank

Question & continued	
Question 8 continued	

Question & continued	Leave
Question 8 continued	
	Q8
	Q <sub>0</sub>
(Total 10 marks)	



**9.** (a) Prove by induction that, for  $n \in \mathbb{N}$ 

$$\sum_{r=1}^{n} r^3 = \frac{1}{4} n^2 (n+1)^2$$

**(5)** 

(b) Using the standard summation formulae, show that

$$\sum_{r=1}^{n} r(r+1)(r-1) = \frac{1}{4} n(n+A)(n+B)(n+C)$$

where A, B and C are constants to be determined.

**(4)** 

(c) Determine the value of n for which

$$3\sum_{r=1}^{n} r(r+1)(r-1) = 17\sum_{r=n}^{2n} r^{2}$$

**(5)** 



	Leave
	blank
Question 9 continued	



Leave blank

Question 9 continued	
yaconon y conomica	

estion 9 continued	

**END**