Please check the examination deta	ils below	before ente	ering your candidate information
Candidate surname			Other names
Pearson Edexcel International Advanced Level	Centre	Number	Candidate Number
Wednesday 7	No	oven	nber 2018
Morning (Time: 2 hours 30 minute	es)	Paper Re	eference WMA02/01
Core Mathemat	ics	C34	
You must have: Mathematical Formulae and Stat	istical 7	Tables (Blu	ue) Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each guestion carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P53789A
©2018 Pearson Education Ltd.
1/1/1/1/1/



- (a) Write $\cos \theta + 4 \sin \theta$ in the form $R \cos (\theta \alpha)$, where R and α are constants, R > 0 and $0 < \alpha < \frac{\pi}{2}$. Give the exact value of R and give the value of α in radians to 3 decimal places.
 - (3)

(b) Hence solve, for $0 \le \theta < \pi$, the equation

$$\cos 2\theta + 4\sin 2\theta = 1.2$$

giving your answers to 2 decimal places.

(5)

Question 1 continued	Leave blank
	Q1
(Total 8 marks)	
(Total & Illal Ks)	



2. A curve C has equation

$$x^3 - 4xy + 2x + 3y^2 - 3 = 0$$

Find an	equation	of the	normal 1	to C at	the	point	(-3,	2),	giving	your	answer	in	the	form
ax + by	c + c = 0	where	a, b and	<i>c</i> are i	integ	gers.								

(7)

Question 2 continued	blank
	Q2
(Total 7 marks)	



Given

 $\cos \theta^{\circ} = p$, where p is a constant and θ° is acute

use standard trigonometric identities to find, in terms of p,

(a) $\sec \theta^{\circ}$

(1)

Leave blank

(b) $\sin(\theta - 90)^{\circ}$

(2)

(c) $\sin 2\theta^{\circ}$

(3)

Write each answer in its simplest form.

6



estion 3 continued	



DO NOT WRITE IN THIS AREA




Figure 1

Figure 1 shows a sketch of part of the curve with equation $y = 8x - xe^{3x}$, $x \ge 0$

The curve meets the x-axis at the origin and cuts the x-axis at the point A.

(a) Find the exact x coordinate of A, giving your answer in its simplest form.

(2)

The curve has a maximum turning point at the point M.

(b) Show, by using calculus, that the x coordinate of M is a solution of

$$x = \frac{1}{3} \ln \left(\frac{8}{1+3x} \right) \tag{5}$$

(c) Use the iterative formula

$$x_{n+1} = \frac{1}{3} \ln \left(\frac{8}{1 + 3x_n} \right)$$

with $x_0 = 0.4$ to calculate the values of x_1 , x_2 and x_3 , giving your answers to 3 decimal places.

(3)

	Le
Question 4 continued	bl
Question 4 continued	



Question 4 continued

	Leave blank
Question 4 continued	
	Q4
(Total 10 marks)	



5.

$$f(x) = \frac{4x^2 + 5x + 3}{(x+2)(1-x)^2} \equiv \frac{A}{(x+2)} + \frac{B}{(1-x)} + \frac{C}{(1-x)^2}$$

(a) Find the values of the constants A, B and C.

(4)

- (b) (i) Hence find $\int f(x) dx$.
 - (ii) Find the exact value of $\int_0^{\frac{1}{2}} f(x) dx$, writing your answer in the form $p + \ln q$, where p and q are constants.

- 4	
-	

uestion 5 continued	



Question 5 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_

Question 5 continued	ы	eave lank
		0.5
		Q5
	(Total 10 marks)	



6. (a) Use binomial expansions to show that, for $|x| < \frac{1}{2}$

$$\sqrt{\frac{1+2x}{1-x}} \approx 1 + \frac{3}{2}x + \frac{3}{8}x^2 \tag{6}$$

(b) Find the exact value of $\sqrt{\frac{1+2x}{1-x}}$ when $x = \frac{1}{10}$

Give your answer in the form $k\sqrt{3}$, where k is a constant to be determined.

(1)

(c) Substitute $x = \frac{1}{10}$ into the expansion given in part (a) and hence find an approximate value for $\sqrt{3}$

Give your answer in the form $\frac{a}{b}$ where a and b are integers. (2)

	Leave
	blank
Question 6 continued	



Question 6 continued

	Leave blank
Question 6 continued	
	06
	Q6
(Total 9 marks)	



7. A curve has equation

$$y = \ln(1 - \cos 2x), \quad x \in \mathbb{R}, 0 < x < \pi$$

Show that

(a)
$$\frac{dy}{dx} = k \cot x$$
, where k is a constant to be found. (4)

Hence find the exact coordinates of the point on the curve where

(b)
$$\frac{\mathrm{d}y}{\mathrm{d}x} = 2\sqrt{3}$$

(4)

astion 7 continued	
estion 7 continued	



Question 7 continued		

Question 7 continued		blank
		07
		Q7
	(Total 8 marks)	



8. (i) Find $\int x \sin x \, dx$

(3)

(ii) (a) Use the substitution $x = \sec \theta$ to show that

$$\int_{1}^{2} \sqrt{1 - \frac{1}{x^{2}}} \, dx = \int_{0}^{\frac{\pi}{3}} \tan^{2}\theta \, d\theta$$
(3)

(b) Hence find the exact value of

$$\int_1^2 \sqrt{1 - \frac{1}{x^2}} \, \mathrm{d}x$$

(4)

		Le bl
Question 8 continued		
	_	



Question 8 continued

Question 8 continued		blank
		Q8
(Tot	tal 10 marks)	



9. A rare species of mammal is being studied. The population *P*, *t* years after the study started, is modelled by the formula

$$P = \frac{900e^{\frac{1}{4}t}}{3e^{\frac{1}{4}t} - 1}, \quad t \in \mathbb{R}, \quad t \geqslant 0$$

Using the model,

(a) calculate the number of mammals at the start of the study,

(1)

(b) calculate the exact value of t when P = 315

Give your answer in the form $a \ln k$, where a and k are integers to be determined.

(4)

(c) (i) Find $\frac{dP}{dt}$

(ii) Hence find the value of $\frac{dP}{dt}$ when t = 8, giving your answer to 2 decimal places.



uestion 9 continued	



Question 9 continued	

Question 9 continued	blank
	_
	_
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	-
	-
	-
	Q9
(Total 9 marks)






Figure 2

Figure 2 shows a sketch of part of the graph with equation y = g(x), where

$$g(x) = \frac{3x - 4}{x - 3}, \quad x \in \mathbb{R}, \quad x < 3$$

The graph cuts the x-axis at the point A and the y-axis at the point B, as shown in Figure 2.

(a) State the range of g.

(1)

- (b) State the coordinates of
 - (i) point A
 - (ii) point B

(2)

(c) Find gg(x) in its simplest form.

(3)

(d) Sketch the graph with equation y = |g(x)|

On your sketch, show the coordinates of each point at which the graph meets or cuts the axes and state the equation of each asymptote.

(3)

(e) Find the exact solution of the equation |g(x)| = 8

(3)



		Leave
		blank
Question 10 continued		
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	
	.	



Question 10 continued

	Leave blank
Question 10 continued	
	Q10
(Total 12 marks)	



11. Relative to a fixed origin O, the line l_1 is given by the equation

$$l_1: \quad \mathbf{r} = \begin{pmatrix} 2\\3\\-1 \end{pmatrix} + \lambda \begin{pmatrix} -1\\4\\3 \end{pmatrix}$$

where λ is a scalar parameter.

The line l_2 passes through the origin and is parallel to l_1

(a) Find a vector equation for l_2

(2)

The point A and the point B both lie on l_1 with parameters $\lambda = 0$ and $\lambda = 3$ respectively.

Write down

- (b) (i) the coordinates of A,
 - (ii) the coordinates of B.

(2)

(c) Find the size of the acute angle between OA and l_1

Give your answer in degrees to one decimal place.

(3)

The point D lies on l_2 such that OABD is a parallelogram.

(d) Find the area of OABD, giving your answer to the nearest whole number.

(3)



	L
uestion 11 continued	ŀ
testion 11 continued	
	1



Leave blank

Question 11 continue	:d		

	Leave
	blank
Question 11 continued	
	011
	Q11
/TT (140 1)	
(Total 10 marks)	



DO NOT WRITE IN THIS AREA




Figure 3

Figure 3 shows a sketch of part of the curve C with parametric equations

$$x = 7t^2 - 5,$$
 $y = t(9 - t^2),$ $t \in \mathbb{R}$

(a) Find an equation of the tangent to C at the point where t = 1

Write your answer in the form ax + by + c = 0, where a, b and c are integers.

(5)

The curve C cuts the x-axis at the points A and B, as shown in Figure 3

- (b) (i) Find the x coordinate of the point A.
 - (ii) Find the x coordinate of the point B.

(3)

The region R, shown shaded in Figure 3, is enclosed by the loop of the curve C.

(c) Use integration to find the area of R.

(5)

	I
uestion 12 continued	'
uosion 12 commucu	



Leave blank

Question 12 continued	

	Leave blank
Question 12 continued	
	0.12
	Q12
(Total 13 marks)	



13. The volume of a spherical balloon of radius rm is Vm³, where $V = \frac{4}{3}\pi r^3$

(a) Find
$$\frac{dV}{dr}$$
 (1)

Given that the volume of the balloon increases with time t seconds according to the formula

$$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{20}{V(0.05t+1)^3}, \qquad t \geqslant 0$$

(b) find an expression in terms of r and t for $\frac{dr}{dt}$ (3)

Given that V = 1 when t = 0

(c) solve the differential equation

$$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{20}{V(0.05t+1)^3}$$

giving your answer in the form $V^2 = f(t)$.

(6)

(d) Hence find the radius of the balloon at time t = 20, giving your answer to 3 significant figures.

(3)

	L
Duration 12 continued	b
Question 13 continued	
	—
	_
	1



Leave blank

Question 13 continued		

	L
Duration 12 continued	b
Question 13 continued	
	—
	_
	1



Question 13 continued	biank
	Q13
(Total 13 marks)	
TOTAL FOR PAPER: 125 MARKS	
END	