

# Risk factors associated with lower extremity stress fractures in runners: a systematic review with meta-analysis

Alexis A Wright,<sup>1</sup> Jeffrey B Taylor,<sup>1</sup> Kevin R Ford,<sup>1</sup> Lindsey Siska,<sup>2</sup> James M Smoliga<sup>1</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2015-094828>).

<sup>1</sup>Department of Physical Therapy, High Point University, High Point, North Carolina, USA

<sup>2</sup>Department of Exercise Science, High Point University, High Point, North Carolina, USA

**Correspondence to**  
Dr Alexis A Wright, High Point University, 833 Montlieu Ave, Drawer 67, High Point, NC 27262, USA;  
[awright@highpoint.edu](mailto:awright@highpoint.edu)

Accepted 26 June 2015

Published Online First  
17 July 2015

## ABSTRACT

**Background** Stress fractures are common overuse injuries with up to 95% occurring in the lower extremities. Among runners, stress fractures account for 15–20% of all musculoskeletal injuries.

**Purpose** We systematically reviewed and critiqued the evidence regarding risk factors associated with increased risk of lower extremity stress fractures in runners.

**Study design** Systematic review.

**Methods** A systematic, computerised literature search of Medline, Embase, PubMed, SPORTDiscus, and CINAHL databases (from database inception through 9 January 2014) using keywords related to risk factors and stress fractures. This systematic review with meta-analysis utilised the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for the search and reporting phases of the study. Inverse variance meta-analyses, using a random effects model were used to summarise ORs.

**Results** 8 articles met the inclusion criteria; 7 were considered low risk. 4 articles qualified for meta-analysis. Results of the meta-analysis identified previous history of stress fracture and female sex as the primary risk factors for future stress fracture with a pooled OR of 4.99 (95% CI 2.91 to 8.56;  $p < 0.001$ ;  $I^2 = 0\%$ ) and 2.31 (95% CI 1.24 to 4.29;  $p < 0.01$ ;  $I^2 = 0\%$ ), respectively.

**Conclusions** Currently, only previous history of stress fracture and female sex are risk factors for lower extremity stress fractures strongly supported by the data.

increased risk of lower extremity stress fractures specifically in runners.

## METHODS

This systematic review was conducted and reported according to the protocol outlined by Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) using a research question developed by PICOS methodology. The PRISMA checklist for this study is available as an online supplementary file. (PICOS is an acronym for population (eg, runners), intervention (eg, risk factors associated with stress fracture), comparison (eg, control group), outcome (eg, OR), and study design (eg, cohort)).

## Identification and selection of the literature

A systematic review was performed to identify risk factors associated with stress fractures in runners. For the purposes of this study, runners are defined as cross-country, track and field, triathlon, duathlon, and recreational running athletes. Articles were identified by searching various databases, including Medline, Embase, PubMed, SPORTDiscus, and CINAHL from database inception through 9 January 2014. Search terms included ('stress fracture' OR 'fatigue fracture' OR 'insufficiency fracture' OR 'march fracture' OR 'overuse fracture' OR 'stress lesions' or 'stress injur' OR 'stress reaction' OR 'bone stress' OR 'bone strain' OR 'repetitive stress injur') AND ('risk' OR 'cohort studies' OR 'mortality'). A hand search was also conducted which included the authors' private collections and the searching of previous related systematic reviews.

## Selection criteria

To qualify for inclusion, studies had to (1) include original data, (2) identify a criterion standard for diagnosis of stress fracture, (3) include an analytical component (ie, the study examined the relationship between risk factor and stress fracture), or sufficient data for these to be calculated, (4) be derived from a prospective cohort of subjects, (5) include lower-extremity stress fractures, (6) be available in full text and (7) written in the English language. Studies were excluded if they did not meet the above inclusion criteria, if they were not specific to the running population, or they failed to include a comparison group of non-injured running athletes.

To identify relevant articles, titles and abstracts of all database-captured citations were independently screened by two reviewers (AAW and JMS) applying the a priori inclusion/exclusion criteria.

## INTRODUCTION

Stress fractures are among the most common overuse injuries reported by active individuals, with up to 95% occurring in the lower extremities.<sup>1</sup> Runners, in particular, are at increased risk with stress fractures accounting for 15–20% of all musculoskeletal-related injuries.<sup>2</sup> Given the many benefits of running and the widespread participation combined with the high incidence of stress fractures, preventing injury among runners has major implications not only for competitive athletes but also for public health.<sup>3</sup>

While it often presumed that risk factors for stress fractures in runners are well known, many recommendations are derived from studies involving military personnel rather than recreational and competitive runners.<sup>4</sup> Additionally, a number of recommendations for prevention of stress fractures in runners are based on cross-sectional studies<sup>5</sup> or theory/opinion. The objective of this study was to conduct a systematic review and meta-analysis of the literature identifying risk factors associated with



CrossMark

**To cite:** Wright AA, Taylor JB, Ford KR, et al. *Br J Sports Med* 2015;**49**:1517–1523.

Full text articles were retrieved if the abstract provided insufficient information to establish eligibility, or if the article passed the first eligibility screening. In situations where disagreement concerning whether to include an article occurred, the two authors discussed the issues and came to a consensus. All criteria were again independently applied by the same two reviewers to the full text of the articles that passed the first eligibility screening. In case of disagreement, a consensus method was used to discuss and solve the disagreement. In the event that agreement could not be reached, a third author (KRF) served as the deciding vote.

### Quality assessment

The two reviewers (AAW and JMS) independently assessed the methodological quality of each study using a risk of bias tool originally derived for measurement of prevalence studies (table 1).<sup>6</sup> The tool includes 10 items to evaluate internal (six items) and external (four items) validity, measurement bias, selection bias and bias related to the analysis (high or low risk), and provides a summary risk of bias assessment based upon the 10 items (low, moderate or high). Item nine, which from the initial risk of bias tool originally dealt with prevalence, was modified to represent the longitudinal nature of risk factor assessment, by characterising studies as high risk if they used a follow-up period of 12 months or shorter. Items were then qualitatively summarised and rated based on an overall assessment of risk bias defined as low (0–3 high-risk items), moderate (4–5 high-risk items), or high risk (6 or greater high-risk items).<sup>7</sup> Disagreements among the reviewers were discussed during a consensus meeting.

### Data extraction

Two reviewers (KRF, JBT) independently extracted information and data regarding study population, study design, patient demographics, reference standard, risk factors and strength of association statistics (eg, OR, adjusted rate ratio (AOR), risk ratio (RR), HR) associated with the risk factors. Reported risk factors only identified by mean differences between groups (ie, p values) were not included.

### Statistical analysis

For studies that provided an OR or RR, calculations were evaluated for accuracy and reported in the analysis. If ORs or RR were not provided, but summary injury data were provided, ratios were calculated by the authors (KRF, JBT) of the current study.<sup>8</sup>

Exploratory meta-analyses were conducted on each risk factor that was described in three or more studies and for which ORs were reported or able to be calculated from included data. Inverse variance meta-analyses, using a random effects model were performed using Comprehensive Meta-Analysis Software (V2.2, Biostat, Englewood, New Jersey, USA). CIs of 95% were calculated, with  $p < 0.05$ , indicating that the true ORs were different than 1.0. Heterogeneity was assessed using an  $I^2$  statistic, with values  $< 25\%$  deemed as low.<sup>9</sup>


### RESULTS

Initially, the search yielded 6957 citations (Medline 1483, Embase 594, PubMed 779, SportDiscus 1700, Cinahl 2401). Of these, 3129 duplicates were deleted, resulting in 3832 titles with abstracts for review. After the first screening, the full texts of 69 potentially eligible citations were retrieved. Following a consensus meeting, a total of eight studies,<sup>10–16</sup> reporting on four independent cohorts, were included in the review (figure 1).

**Table 1** Methodological quality assessment scores of included articles

Article	External validity				Internal validity			The length of the assessment period was appropriate	The numerator and denominator of the parameter of interest was reported	Overall score
	Population was a close representation of national population	Sampling frame was a close representation of population	Randomised sample or census data were used	Minimal non-response bias	Data collected directly from the subjects	Acceptable case definition was used	Instrumentation was reliable and valid	The same mode of data collection was used for all subjects		
Bennell <i>et al</i> <sup>10</sup>	High	Low	High	Low	Low	Low	Low	Low	High	Low
Bennell <i>et al</i> <sup>12</sup>	High	Low	High	Low	Low	Low	Low	Low	High	Low
Bennell <i>et al</i> <sup>11</sup>	High	Low	High	Low	Low	Low	Low	Low	High	Low
Cobb, 2007 <sup>17</sup>	Low	Low	High	Low	Low	High	Low	High	Low	Low
Kelsey <i>et al</i> <sup>13</sup>	Low	Low	High	Low	Low	High	Low	Low	Low	Low
Nieves <i>et al</i> <sup>14</sup>	Low	Low	High	Low	Low	High	Low	Low	Low	Low
Tenforde <i>et al</i> <sup>15</sup>	High	Low	Low	Low	High	High	High	Low	Low	Moderate
Yagi <i>et al</i> <sup>16</sup>	High	Low	Low	Low	High	Low	Low	Low	High	Low

**Figure 1** Flow diagram of the literature screening process.



The results of the quality assessment are shown in table 1. Of the eight included studies, seven<sup>10–14 16</sup> were classified as low risk, and one study<sup>15</sup> was classified as moderate risk. Most methodological shortcomings concerned the following items: the study's target population was clearly not representative of the national population, a random sample was not used, data was collected via a proxy (eg, questionnaire) as opposed to directly from the subject, and failure to calculate appropriate statistics (eg, OR).

A summary of the individual study characteristics is presented in the online supplementary table. Three studies<sup>13–15</sup> investigated nutritional status/dietary habits, four studies<sup>10 13 15 17</sup> investigated use of oral contraceptives, two studies investigated training regimen,<sup>10 15</sup> three studies<sup>12 13 15</sup> investigated history of previous fractures, three studies<sup>12 15 16</sup> investigated sex, three studies<sup>10 13 15</sup> investigated menstrual status, and three studies<sup>11 13 15</sup> investigated body composition and bone density. The number of patients with follow-up ranged from 95<sup>12</sup> to 701.<sup>15</sup> The mean follow-up duration ranged from 11.4 months<sup>15</sup> to 36<sup>16</sup> months. A positive stress fracture diagnosis was based on four different criteria: conventional radiograph, MRI, nuclear scintigraphy, or CT.

### Risk factors and associations with stress fracture

#### Extrinsic variables

##### Nutrition/dietary habits


Three studies reported on dietary intake and eating behaviours with conflicting findings.<sup>13–15</sup> Nieves<sup>14</sup> reported higher consumption of calcium (HR 0.53, 95% CI 0.29 to 0.97), servings of dairy product per day (HR 0.60, 95% CI 0.40 to 0.89), skim

milk (HR 0.38, 95% CI 0.16 to 0.90), total milk consumption (HR 0.43, 95% CI 0.20 to 0.89), and high dairy/low fat dietary patterns (HR 0.32, 95% CI 0.10 to 0.96) were related to a significantly reduced rate of stress fracture (see online supplementary table). Kelsey *et al*<sup>13</sup> found that lower average daily servings of dairy products (per one serving decrease) were related to an increased rate of stress fracture (RR 1.41, 95% CI 1.01 to 1.96). Interestingly, when reported as calcium intake per 100 mg decrease, Kelsey *et al*<sup>13</sup> found no significant association between calcium intake and decreased or increased rate of stress fractures. Tenforde *et al*<sup>15</sup> found that female athletes who used a calcium supplement were at 3 times higher risk of stress fracture (OR 3.36, 95% CI 1.4 to 7.9). The same was not true for male athletes (OR 0.90, 95% CI 0.19 to 4.28). Limitations regarding study protocol were questioned as athletes were asked late in the study regarding calcium intake and calcium supplementation was also strongly correlated to previous history of stress fracture. Therefore, injured athletes may have been prescribed calcium supplementation as part of their treatment plan versus primary prevention. History of diagnosed eating disorder (anorexia nervosa or bulimia nervosa) was reported in a single study<sup>15</sup> with a significant association with increased risk of stress fracture (OR 259.8, 95% CI 30.8 to 2193.40).

##### Use of oral contraceptives

Four studies<sup>10 13 15 17</sup> reported on the use of contraceptives among female athletes. In all four studies, use or non-use of an oral contraceptive was not significantly related to a reduced or

**Figure 2** Pooled OR with 95% CI for previous history of stress fracture. Kelsey *et al*.<sup>13</sup> \*OR calculated from adjusted rate ratio.



increased rate of stress fracture with ORs ranging from 0.55 to 2.22 (see online supplementary table).

### Training regimen

Two studies<sup>10 15</sup> reported on training regimen as a risk factor. Tenforde *et al*<sup>15</sup> defined training regimen as the average weekly running distance in the past year. Athletes running >32 km/week were found to be at 2 times higher risk of injury (OR 2.40; 95% CI 1.20 to 4.82). Females, in particular, running >32 km/week were at 3 times increased risk of stress fracture as compared with those running <32 km/week (OR 3.08, 95% CI 1.31 to 7.20). Bennell *et al*<sup>10</sup> found no significant associations between premenarcheal training and increased risk of stress fracture.

### Intrinsic variables

#### Previous history of stress fracture

Three studies<sup>12 13 15</sup> reported on previous history of stress fracture and its association with increased risk of future stress fracture. All three studies had similar findings in that athletes with a previous history of fracture were at increased risk of developing a future stress fracture with ORs ranging from 2.90 to 6.36. An exploratory meta-analysis confirmed the individual study results with runners with a previous history of stress fracture at 5 times higher risk of a future stress fracture (figure 2, table 2; OR 4.99; 95% CI 2.91 to 8.56;  $p<0.001$ ;  $I^2=0\%$ ).

### Sex

Three studies reported on sex and its association with increased risk of stress fracture with conflicting findings.<sup>12 15 16</sup> Yagi *et al*<sup>16</sup> found an increased risk of stress fractures among females (OR 3.10, 95% CI 1.20 to 8.00). Specifically, 5% of males compared to 15% of females sustained a stress fracture during the prospective study of high school running teams. By contrast, Bennell *et al*<sup>12</sup> (OR 1.08, 95% CI 0.40 to 2.91) and Tenforde *et al*<sup>15</sup> (OR 1.35, 95% CI 0.65 to 2.82) found no significant association between sex and increased risk of stress fracture. An exploratory meta-analysis with study subgroups focused specifically on distance runners was performed based on the increased risk of stress fracture in females compared with males (figure 3,

table 2). Females were at 2.3 times higher risk compared with males (figure 3; OR 2.31; 95% CI 1.24 to 4.29;  $p=0.008$ ;  $I^2=0\%$ ).

### Menstrual characteristics

Three studies<sup>10 13 15</sup> reported on menstrual characteristics for female athletes defined in terms of age at menarche, amenorrhoeic, or oligomenorrhoeic with conflicting findings. Bennell *et al*<sup>10</sup> found no significant association between female athletes currently defined as oligomenorrhoeic or amenorrhoeic and increased risk of stress fracture (OR 3.43, 95% CI 0.62 to 18.88). Past history of oligomenorrhoea (OR 1.32, 95% CI 0.29 to 5.99) or amenorrhoea (OR 2.60, 95% CI 0.62 to 10.96) was also found not to be significantly associated with increased risk of stress fracture.<sup>10</sup> Kelsey *et al*<sup>13</sup> also reported no association between history of menstrual irregularity and increased risk of stress fracture (RR 1.90, 95% CI 0.66 to 5.51). By contrast, Tenforde *et al*<sup>15</sup> found that female athletes with later age of menarche ( $\geq 15$  years) and currently amenorrhoeic were at five times (OR 5.30, 95% CI 2.11 to 13.34) and nearly three times (OR 2.73, 95% CI 1.16 to 6.4) increased risk of stress fractures, respectively.

### Body composition and bone density

Two studies<sup>11 13</sup> reported on varying aspects of bone mineral content defined as total bone mineral content, bone mineral density, and whole-body skeletal area and their association with increased risk of stress fracture. Neither of these studies provided enough information to calculate an OR. Kelsey *et al*<sup>13</sup> found significantly lower values for whole-body bone mineral content (RR 1.79, 95% CI 1.02 to 3.13), spine bone mineral density (RR 1.89, 95% CI 1.04 to 3.45) and whole body skeletal area (RR 1.89, 95% CI 1.06 to 3.33) in women who developed stress fractures (see online supplementary table). Kelsey *et al*<sup>13</sup> also reported a significant association between lean body mass (per kg decrease) and increased risk of stress fracture (OR 1.14, 95% CI 1.01 to 1.28). Tenforde *et al*<sup>15</sup> found a significant association between BMI<19 and increased risk of stress fracture (OR 2.76, 95% CI 1.18 to 6.46).

## DISCUSSION

Our systematic review aimed to identify risk factors associated with increased risk of lower extremity stress fractures in the running population. In this review of eight studies, there was a striking lack of conclusive evidence for any risk factor, but with two exceptions: previous history of stress fracture and female sex-predicted increased risk of future stress fracture.


Although history of a previous stress fracture is an intrinsic, non-modifiable risk factor in terms of prevention, this finding stresses the importance of identification of modifiable risk

**Table 2** Pooled estimates from meta-analysis

Risk factor	Number studies sample size (N)	OR (95% CI)	Z value	p Value
History of stress fracture	3 (N=939) <sup>12 13 15</sup>	4.99 (2.91 to 8.56)	5.84	<0.01
Female sex	3 (N=480) <sup>12 15 16</sup>	2.31 (1.24 to 4.29)	2.65	0.01



**Figure 3** Pooled OR with 95% CI for female sex. Bennell *et al.*<sup>12</sup> ≥OR calculated from male/female distance runners only. Tenforde *et al.*<sup>15</sup> ‡OR calculated from male/female runners >32 km/week.



factors responsible for the initial injury and subsequent effective rehabilitation to reduce risk of future stress fracture. The deficient evidence highlighted in this systematic review identifies the need for further research.<sup>17</sup>

### Why is the evidence scant?

It has previously been acknowledged that prospective studies examining stress fractures are difficult to perform given the large populations that must be followed for long time periods, and thus, retrospective studies are often pursued, even though they are unable to establish a causal relationship.<sup>18</sup> Additionally, one could argue that the lack of available evidence may be secondary to the complex nature of stress fracture research in the running population. One of the complications in stress fracture research in runners is the interdependence of variables and how individual differences may influence response to confounding factors. Tenforde *et al.*'s<sup>15</sup> findings, in which some apparently significant risk factors (eg, training volume, body mass, racing performance) became non-significant when multivariate analysis was conducted, serves as a prime example of the interdependent nature of risk factors for stress fractures. While many of these factors were examined in various studies that met the inclusion criteria, they were often not examined synchronously. The individual nature of stress fracture risk, and thus, the need to identify risk factors to prevent them, is also clear when one considers athletes performing nearly identical training regimens within a team. While some athletes remain injury free, others develop soft tissue injuries (eg, tendinopathy), whereas others develop stress fractures at different sites. Finally, there is emerging evidence that various single nucleotide polymorphisms may be associated with stress fractures in athletes,<sup>19</sup> however, considerable research is necessary before more definitive conclusions can be made regarding genetic risks for stress fractures in runners.

### Bone mineral density and calcium as risk factors

Mechanical loading is associated with increased bone mineral density,<sup>20</sup> thus, runners with greater training volumes should, theoretically, have higher bone mineral density. Although ORs could not be computed from mean data, bone density in runners with a history of stress fracture was lower than those without stress fractures, yet greater than that of non-athletes.<sup>10</sup> This suggests that healthy bone mineral density requirements of active running individuals may be greater than that required by the general public.

This systematic review found conflicting results regarding whether calcium supplementation is associated with increased<sup>15</sup> or decreased<sup>13,14</sup> risk for stress fractures. Though dietary interventions focused on calcium intake are theoretically beneficial

to reduce risk of stress fracture in runners, no prospective clinical trial data is available to support this recommendation. A full understanding of dietary risk factors is further complicated by differences in methodology used to assess nutrition, as well as variety and crossover in predictor variables (eg, dairy intake vs calcium intake).

### Menstrual dysfunction

Although menstrual dysfunction is commonly cited as a risk factor for stress fractures in athletes, Tenforde *et al.*'s<sup>15</sup> was the only study identified which described an association between the variables, in that age of menarche and current amenorrhoea were associated with significant increases in risk for stress fracture. Conflicting findings in this realm may relate to fallacies within the traditional 'female athlete triad' paradigm.<sup>21,22</sup> Menstrual dysfunction may be considered simply as one group of signs which occurs secondary to the issue of 'relative energy deficit in sports (RED-S)'.<sup>21</sup> It is possible that female athletes who experience RED-S, and are at elevated risk of stress fracture, have suboptimal menstrual function that is not clinically apparent,<sup>21,22</sup> and such subclinical issues are not recognised in risk factor studies.

### Biomechanical risk factors

Biomechanical factors, including kinetic, kinematic and static alignment variables, are commonly implicated as a risk factor for stress fractures, however, no studies which met the inclusion criteria examined biomechanical factors. One commonly cited study prospectively links biomechanical variables (average and instantaneous vertical loading rate, tibial shock) to stress fractures in competitive female runners, however, it is not a peer-reviewed article, but rather a 2004 research abstract with only five individuals with stress fractures.<sup>23</sup> Other studies which have focused on biomechanical 'predictors' in runners have chiefly examined biomechanical differences (ie, peak hip adduction, knee rotation, rearfoot eversion angle) in individuals with a previous history of stress fracture,<sup>24–27</sup> or have examined biomechanical variables which, theoretically, could lead to stress fracture without prospectively tracking the incidence of injury.<sup>28–30</sup> Also noticeable by its absence is the almost complete lack of investigation of the association of biomechanically relevant clinical examination components with stress fracture risk such as arch height, alignment of lower extremity, range of motion, and leg-length inequality.

### Military population data should not be translated directly to runners

A wealth of research has been conducted on risk factors associated with stress fractures; however, much of this data comes

from studies involving military personnel, with far more sampling populations and considerably larger sample sizes than those specific to runners.<sup>4</sup> Studies involving military personnel include smokers and individuals with poor overall physical fitness, both of whom would not be expected in a population of serious runners, yet are significantly associated with stress fractures<sup>31</sup> and overall training-related injuries.<sup>32–33</sup> Military recruits often run <10 miles per week<sup>34</sup> and often take 12 weeks to accumulate 40 miles of running,<sup>35</sup> whereas runners often perform 40 miles of running at various intensities within 1 week, with some competitive runners exceeding 100 miles per week.<sup>36</sup> Additionally, soldiers regularly carry heavy loads<sup>37</sup> and use footwear that is not as effective for injury prevention as typically found in running shoes,<sup>38</sup> further prohibiting a direct comparison with running athletes. Stress fractures in the military often occur within the first 16 weeks of training,<sup>4</sup> but risk factors during this initial acclimation period are not likely applicable to athletes who experience their first stress fracture after years of running. Although military data are commonly extrapolated to the general running population, this is likely inappropriate due to major differences between these populations.

### Clinical implications

We found little or conflicting evidence for extrinsic or modifiable risk factors such as training regimen, nutritional status, or biomechanics. Only one study reported on training volume (eg, mileage) associated with increased risk of stress fracture, however, the reported threshold for increased risk was a relatively low volume (32 km/week) in relation to the training regimens many runners undertake.<sup>36–39</sup> Additionally, running volume itself is not an overall indicator of mechanical stress placed upon the musculoskeletal system. A typical training week may involve a combination of different intensities of running, which place different magnitudes of load on the skeleton. Rapid changes in training volume and intensity are often considered in the development of injuries, though this is difficult to quantify in epidemiological studies.

Thus, future research examining training factors should also consider incorporating a holistic measure, such as training impulse,<sup>40</sup> while also seeking to describe the stability of training practices, and reporting injury rates in terms of both time (eg, injuries per week) and volume (eg, injuries per mile). In the absence of methodologically sound high-quality studies which quantify training practices, including volume, intensity and progression, clinicians have little evidence to support the widespread anecdote that higher mileages are associated with increased risk of stress fracture, or the corresponding recommendation that reducing running volume can prevent stress fractures.

### Limitations

There is an overall lack of prospective studies focused on stress fractures within the running population, some of which do not include the OR associated with the identified risk factor which is recognised as the proper method for reporting risk factor ratios. Only four independent cohorts made up the eight included studies of this review highlighting the complete lack of research performed in the running population. While three of the cohorts exclusively focused on middle and long distance runners, one cohort included a small sample of track and field athletes from other events.<sup>12</sup> Likewise, this systematic review focused on lower extremity stress fractures, but some cohorts included isolated cases of pelvic stress fractures. Studies which examined stress fracture risk in a heterogeneous group of

athletes,<sup>41</sup> or those which examined general risk factors for all types of running injuries<sup>42</sup> were not included, as data reported in these broader studies are not sufficient to determine the risk factors specific to stress fracture in runners.

The available data may be at risk of bias given the considerable variation among the studies with respect to diagnostic criteria (radiograph, nuclear scintigraphy, or MRI), with studies using scintigraphy and MRI potentially reporting a higher incidence. With only a single cohort<sup>43</sup> reporting distribution of stress fracture by site and mode of diagnosis, risk factors for stress fractures at specific anatomic sites (eg, fifth metatarsal vs femoral neck) could not be considered, and this may be important if the mechanism of injury varies by location.

In conclusion, our systematic review revealed little evidence for most risk factors for stress fracture within the running population. Previous stress fracture was identified as the greatest risk factor for a stress fracture, however, it is unknown whether this represents new injuries or a previous stress fracture which was not fully healed. Female runners appear to be at greater risk of stress fractures than their male counterparts, and the evidence is inconclusive as to whether this risk may be elevated in females with a history of menstrual dysfunction, low bone mineral density, low calcium intake, and training at higher volumes.

Given the current state of literature, perhaps the most important role for clinicians to reduce the impact of stress fractures are through proper education and time-efficient diagnosis. When

### What are the new findings

- ▶ Only eight studies, composed of four unique data sets, have specifically examined stress fracture risk factors in runners.
- ▶ Runners at greatest risk of stress fracture are those with a previous history of stress fracture and females.
  - There are conflicting findings regarding nutritional factors/dietary history and menstrual status/history as risk factors for stress fractures in runners.
  - There is no evidence that use of oral contraceptives is related to stress fracture risk in runners.
  - One study found that running >32 km (>20 miles) per week was a risk factor for stress fracture, however, this relatively low volume of training commonly was exceeded by most runners.
- ▶ Future research should investigate how running-specific training practices, nutritional factors, biomechanics and other factors influence the risk of stress fractures in runners.

### How might it impact on clinical practice in the near future

- ▶ Female runners with a history of a stress fracture in combination with clinical findings suggestive of a lower extremity stress fracture should warrant early referral for diagnostic imaging.
- ▶ Surprisingly, there was no evidence to suggest training practices (eg, surface, terrain, volume) is associated with increased risk of stress fracture in runners.
- ▶ Clinicians are encouraged to use caution in extrapolating risk factor findings from military studies to the running population given the major differences between populations.

patients present to clinicians for pre-participation physicals prior to team sport enrolment or beginning a self-guided training programme, they should be encouraged to beware of the signs of musculoskeletal injuries and to take appropriate measures at the onset of suspected injury.

**Contributors** The authors declare that all named authors made substantial contributions to the conception or design of the work; drafting and revising the work, final approval of the manuscript to be published, and are in agreement to be accountable for all aspects of the work.

**Competing interests** None declared.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- Liong SY, Whitehouse RW. Lower extremity and pelvic stress fractures in athletes. *Br J Radiol* 2012;85:1148–56.
- Barnes A, Wheat J, Milner C. Association between foot type and tibial stress injuries: a systematic review. *Br J Sports Med* 2008;42:93–8.
- Engelbrechtsen L, Bahr R, Cook JL, et al. The IOC Centres of Excellence bring prevention to Sports Medicine. *Br J Sports Med* 2014;48:1270–5.
- Wentz L, Liu PY, Haymes E, et al. Females have a greater incidence of stress fractures than males in both military and athletic populations: a systemic review. *Mil Med* 2011;176:420–30.
- Duckham RL, Peirce N, Meyer C, et al. Risk factors for stress fracture in female endurance athletes: a cross-sectional study. *BMJ Open* 2012;2:e001920.
- Hoy D, Bain C, Williams G, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum* 2012;64:2028–37.
- Cook CE, Taylor J, Wright A, et al. Risk factors for first time incidence sciatica: a systematic review. *Physiother Res Int* 2014;19:65–78.
- Portney LG, Watkins MP. *Foundations of Clinical Research: Applications to practice*. 3rd edn. Upper Saddle River, NJ: Prentice Hall, 2008.
- Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- Bennell KL, Malcolm SA, Thomas SA, et al. Risk factors for stress fractures in track and field athletes: a twelve-month prospective study. *Am J Sports Med* 1996;24:810–18.
- Bennell KL, Malcolm SA, Brukner PD, et al. A 12-month prospective study of the relationship between stress fractures and bone turnover in athletes. *Calcif Tissue Int* 1998;63:80–5.
- Bennell KL, Malcolm SA, Thomas SA, et al. The incidence and distribution of stress fractures in competitive track and field athletes. A twelve-month prospective study. *Am J Sports Med* 1996;24:211–17.
- Kelsey JL, Bachrach LK, Procter-Gray E, et al. Risk factors for stress fracture among young female cross-country runners. *Med Sci Sports Exerc* 2007;39:1457–63.
- Nieves JW, Melsop K, Curtis M, et al. Nutritional factors that influence change in bone density and stress fracture risk among young female cross-country runners. *PM R* 2010;2:740–50; quiz 94.
- Tenforde AS, Sayres LC, McCurdy ML, et al. Identifying sex-specific risk factors for stress fractures in adolescent runners. *Med Sci Sports Exerc* 2013;45:1843–51.
- Yagi S, Muneta T, Sekiya I. Incidence and risk factors for medial tibial stress syndrome and tibial stress fracture in high school runners. *Knee Surg Sports Traumatol Arthrosc* 2013;21:556–63.
- Cobb KL, Bachrach LK, Sowers M, et al. The effect of oral contraceptives on bone mass and stress fractures in female runners. *Med Sci Sports Exerc* 2007;39:1464–73.
- Lohmander LS, Roos EM. The evidence base for orthopaedics and sports medicine. *BMJ* 2015;350:g7835.
- Warden SJ, Davis IS, Fredericson M. Management and prevention of bone stress injuries in long-distance runners. *J Orthop Sports Phys Ther* 2014;44:749–65.
- Varley I, Hughes DC, Greeves JP, et al. RANK/RANKL/OPG pathway: genetic associations with stress fracture period prevalence in elite athletes. *Bone* 2015;71:131–6.
- Ishikawa S, Kim Y, Kang M, et al. Effects of weight-bearing exercise on bone health in girls: a meta-analysis. *Sports Med* 2013;43:875–92.
- Mountjoy M, Sundgot-Borgen J, Burke L, et al. The IOC consensus statement: beyond the Female Athlete Triad—Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med* 2014;48:491–7.
- Mountjoy M, Sundgot-Borgen J, Burke L, et al. Authors' 2015 additions to the IOC consensus statement: Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med* 2015;49:417–20.
- Davis IS, Milner C, Hamill J. Does increased loading during running lead to tibial stress fractures? A prospective study. *Med Sci Sports Exerc* 2004;36:S58.
- Milner CE, Davis IS, Hamill J. Free moment as a predictor of tibial stress fracture in distance runners. *J Biomech* 2006;39:2819–25.
- Milner CE, Hamill J, Davis IS. Distinct hip and rearfoot kinematics in female runners with a history of tibial stress fracture. *J Orthop Sports Phys Ther* 2010;40:59–66.
- Milner CE, Ferber R, Pollard CD, et al. Biomechanical factors associated with tibial stress fracture in female runners. *Med Sci Sports Exerc* 2006;38:323–8.
- Pohl MB, Mullineaux DR, Milner CE, et al. Biomechanical predictors of retrospective tibial stress fractures in runners. *J Biomech* 2008;41:1160–5.
- Clansey AC, Hanlon M, Wallace ES, et al. Effects of fatigue on running mechanics associated with tibial stress fracture risk. *Med Sci Sports Exerc* 2012;44:1917–23.
- Mizrahi J, Verbitsky O, Isakov E. Shock accelerations and attenuation in downhill and level running. *Clin Biomech (Bristol, Avon)* 2000;15:15–20.
- Milgrom C, Finestone A, Segev S, et al. Are overground or treadmill runners more likely to sustain tibial stress fracture? *Br J Sports Med* 2003;37:160–3.
- Cowan DN, Bedno SA, Urban N, et al. Step test performance and risk of stress fractures among female army trainees. *Am J Prev Med* 2012;42:620–4.
- Jones BH, Cowan DN, Tomlinson JP, et al. Epidemiology of injuries associated with physical training among young men in the army. *Med Sci Sports Exerc* 1993;25:197–203.
- Knapik JJ, Sharp MA, Canham-Chervak M, et al. Risk factors for training-related injuries among men and women in basic combat training. *Med Sci Sports Exerc* 2001;33:946–54.
- Trank TV, Ryman DH, Minagawa RY, et al. Running mileage, movement mileage, and fitness in male U.S. Navy recruits. *Med Sci Sports Exerc* 2001;33:1033–8.
- Knapik JJ, Trone DW, Swedler DL, et al. Injury reduction effectiveness of assigning running shoes based on plantar shape in Marine Corps basic training. *Am J Sports Med* 2010;38:1759–67.
- Spillsbury KL, Fudge BW, Ingham SA, et al. Tapering strategies in elite British endurance runners. *Eur J Sport Sci* 2014:1–7.
- Knapik JJ, Reynolds KL, Harman E. Soldier load carriage: historical, physiological, biomechanical, and medical aspects. *Mil Med* 2004;169:45–56.
- House C, Reece A, Roiz de Sa D. Shock-absorbing insoles reduce the incidence of lower limb overuse injuries sustained during Royal Marine training. *Mil Med* 2013;178:683–9.
- Rauh MJ. Summer training factors and risk of musculoskeletal injury among high school cross-country runners. *J Orthop Sports Phys Ther* 2014;44:793–804.
- Borresen J, Lambert MI. The quantification of training load, the training response and the effect on performance. *Sports Med* 2009;39:779–95.
- Korpelainen R, Orava S, Karpakka J, et al. Risk factors for recurrent stress fractures in athletes. *Am J Sports Med* 2001;29:304–10.
- Saragiotto BT, Yamato TP, Hespanhol Junior LC, et al. What are the main risk factors for running-related injuries? *Sports Med* 2014;44:1153–63.