

Rapid #: -15586379

CROSS REF ID: **1012122**

LENDER: **HUA :: Main Library**

BORROWER: **COD :: Main Library**

TYPE: Article CC:CCL

JOURNAL TITLE: Australian family physician

USER JOURNAL TITLE: Australian family physician

ARTICLE TITLE: Stress fractures of the lower limbs in runners

ARTICLE AUTHOR: Fitch, K D

VOLUME: 13

ISSUE: 7

MONTH: 07

YEAR: 1984

PAGES: 511-

ISSN: 0300-8495

OCLC #:

Processed by RapidX: 1/1/2020 10:06:06 PM



This material may be protected by copyright law (Title 17 U.S. Code)

RapidX Upload

14 **Rapid #: -15586379**

Status	Rapid Code	Branch Name	Start Date
New	COD	Main Library	01/01/2020 07:58 AM
Pending	HUA	Main Library	01/01/2020 07:59 AM
Batch Printed	HUA	Main Library	01/01/2020 08:05 AM

CALL #: MS 610 A93 F19 P
LOCATION: HUA :: Main Library :: medse

REQUEST TYPE: Article CC:CCL
JOURNAL TITLE: Australian family physician
USER JOURNAL TITLE: Australian family physician
HUA CATALOG TITLE: Australian family physician
ARTICLE TITLE: Stress fractures of the lower limbs in runners
ARTICLE AUTHOR: Fitch, K D
VOLUME: 13
ISSUE: 7
MONTH: 07
YEAR: 1984
PAGES: 511- 515
ISSN: 0300-8495
OCLC #: HUA OCLC #: 5918923
CROSS REFERENCE ID: [TN:1012122][ODYSSEY:206.107.44.246/COD]
VERIFIED:

BORROWER: COD :: Main Library



This material may be protected by copyright law (Title 17 U.S. Code)
1/2/2020 8:57:36 AM

香港大學
圖書館



THE UNIVERSITY OF HONG KONG LIBRARIES
Lending Services Division
Pokfulam Road, Hong Kong

Circulation enquires: (852)39175945 / 39172202

E-mail: libcir@hku.hk

Interlibrary Loan enquiries: (852)3917-2215 / 3917-5895

E-mail: interlib@hku.hk

Fax: (852)25174615

Associate Librarian, Engagement and Lending Services: 39177001

University of Hong Kong Libraries

© The copy is for purposes of private study or scholarly research only.

You should delete the file as soon as a single paper copy has been printed out satisfactorily.

Stress fractures of the lower limbs in runners

K D Fitch

Stress fractures are a common sports injury and running can cause fractures in most bones from the metatarsals to the pubic rami. Pain after an increase in or modification to a running programme is the initial symptom; there will be point tenderness over bone. Rest from running for six to 10 weeks is necessary but alternative exercise must be prescribed for those 'addicted' to running.

The first description (in 1855) of a stress fracture is accredited to Breihaupt, a Prussian army doctor, who reported painful swollen feet in soldiers after long marches.¹ Stechow described metatarsal fractures in soldiers in 1897² and the term 'march fractures' has endured. Four decades elapsed before stress fractures were noted in athletes (again in the German literature).³ Devas, a London orthopaedic surgeon, has been the most prolific author on the subject and his excellent monograph is recommended.⁴



Prepared by request

K D Fitch, MD, is physician to the Sports Injury Clinic at the Royal Perth Rehabilitation Hospital, national convenor of the elective in sports medicine and Senior Medical Director of the 1984 Australian Olympic Team.

The exponential increase in the number of joggers, 'fun' and marathon runners over the past decade has been associated with an 'epidemic' of stress fractures in the bones of the lower limbs. Stress fractures are not confined to runners or the legs and occur in the lumbar pars interarticularis of gymnasts⁵ and fast bowlers; the ribs of weight lifters;⁴ the humerus of throwers and arm wrestlers;⁶ and the hook of the hamate in golfers and tennis players.⁷

Recreational athletes attend their family physician with symptoms of stress fractures but they frequently remain undiagnosed.

Causes

Causes of stress fractures in runners are listed in Table 1.

In many instances there is a combination of factors but the most common is a sudden increase in the training distance. Long distance runs are essential in the preparation for a marathon (42 km) and the more experienced the runner the more likely is a training error the cause of the fracture.

Pathophysiology

The underlying factor in the accelerated remodelling of bone (the consequence of stress overload)^{4, 8} is muscular forces acting on bone. Osteoclastic activity commences the remodelling process and results in resorption, rarefaction and, if the stress is continued, microfractures. With rest or reduced activity osteoblasts lay down a new and stronger bone matrix better able to withstand stress; continued running will convert a microfracture to a macrofracture. Healing involves endosteal and periosteal callus formation to buttress the cortex and focal sclerosis and these are its radiological features.⁴

Clinical features

The onset of a stress fracture can be sudden or insidious; dramatic development while running is uncommon and often preceded by minor symptoms. Usually the fracture begins gradually and is associated with a history of a sudden increase in the amount, frequency or intensity of training. The first symptom is pain after a run which is relieved rapidly by rest. Perseverance with running will increase the severity of pain and its resolution will require a longer period of rest. The second stage is pain as the run begins, often ceasing after a short distance but returning before the run is finished; night pain now might prevent sleep. Finally the pain becomes constant and prevents running.

Examination reveals point tenderness over bone at a typical stress fracture site. There might be slight swelling but a full and pain free range

Table 1
Causative factors in runners' stress fractures

Training errors

Sudden increase in the amount, frequency or intensity of training

Biomechanical factors

Pronated or cavus feet; fore or rear foot varus; tibial varum; discrepancy in leg length

Surfaces and terrain

Change of running surface (eg grass to road); running down or across the face of hills

Running mechanics

Gait is faulty or becomes so when fatigued

Footwear

Shoes damaged or inappropriate for the feet or running surface

Temporary cessation of training

Training resumed after inactivity, especially if previous levels attempted immediately




Figure 1. Right: compression stress fracture in calcaneus of 45-year-old male after three weeks of recreational jogging on footpath.



Figure 3. Bilateral stress fractures of the shafts of the second metatarsal in a 16-year-old boy.




Figure 2. Compression stress fracture of the medial tibial plateau in a 37-year-old jogger.

of passive movements of the proximal and distal joints is customary (except with fractures close to the hip joint). In the majority, resisted contraction of the muscles spanning the injured region of bone is painless.

Type and site

Devas classified stress fractures radiologically as: distraction (oblique, longitudinal, transverse); and compression.⁴

The tibia is a classical site of oblique distraction fractures. Distinction between fracture lines and nutrient arteries can be difficult but fracture lines tend to be 'down and out', that is extend distally from the inner rim to the outer cortex.⁴ A transverse fracture has the potential to extend into a complete fracture which can cause displacement (for example neck of femur).

Compression fractures in which no fracture line is visible are typified in the calcaneus (Figure 1) and the medial tibial plateau (Figure 2). The clinical features are diagnostic; X-ray after four or five weeks will show an area of sclerosis.

Bilateral fractures are not uncommon in the tibiae, fibulae, metatarsals, femora and calcanei (Figure 3). Mul-

tiple fractures in the same bone (especially the tibia) are encountered occasionally and confirm that the healed region of a fully united fracture is able to withstand stress better than the rest of the bone.

Sites of stress fractures are predictable, as exemplified in the metatarsals where fractures of the shafts and neck region of the second and third are most common (Figure 3). The first and fifth metatarsals are overloaded less often than the fourth. In the tarsus, the calcaneus and navicular are the bones affected. Joggers fracture the calcaneus; sprinters, hurdlers and high jumpers fracture the navicular. Navicular fractures are difficult to diagnose radiologically and often result in non union and the need

Stress fractures of the fibula occur five to eight centimetres proximal to the tip of the malleolus (Figure 4). Fractures are seen at most levels of the tibia but usually the upper third in adolescents (Figure 5) and the middle and lower thirds and medial plateau in adults (Figure 2).

- Joggers fracture the calcaneus; sprinters fracture the navicular.

Erythrocin[®]
When the defence system falls short,
the body needs an ally.



When the signs point to oestrogen excess...

Breast tenderness →

Dysmenorrhoea →

Nausea →

Migraine →

← Nordette

Nordette.* The more gentle way.

No combined low dose oral contraceptive contains less oestrogen than Nordette.

Perhaps that's why it's so well received by women whose bodies are happier with a little less oestrogen . . . and why it continues to be the most popular O.C. formulation world-wide.

Each active tablet contains 150mcg levonorgestrel, 30mcg ethinyloestradiol.
N.H.S. General Benefit: 2 cycle packs, 2 repeats, 21 and 28 day packs.

*trademark



Wyeth Pharmaceuticals Pty. Limited, incorporated in N.S.W.,
Gregory Place, Parramatta, N.S.W.

WYE001HAMMOND



Figure 4. Typical site and appearance of a fibula stress fracture.

Femoral stress fracture can develop in the region of insertion of adductor magnus, longus and brevis and in the femoral neck. Failure to diagnose stress fracture of the neck of femur (or the patient's persistence with running despite contrary advice) can result in a complete fracture with displacement which requires internal fixation.⁹ Stress fracture of the inferior pubic ramus¹⁰ (Figure 6), which is slow to heal, is a differential diagnosis of a femoral neck fracture.

Diagnosis

It is essential to consider stress frac-

- Stress fracture should be considered in the differential diagnosis of bone pain.

ture as a diagnostic possibility in any patient presenting with bone pain.

Confirmation of the diagnosis can be obtained by bone scan, conventional radiography, tomography and computerised axial tomography (CAT scan). A radionuclide bone scan using 99 m technetium labelled diphosphonate will be positive almost immediately;¹¹ it is valuable when the clinical diagnosis is in doubt (for example navicular fractures; Figure 7) or evidence is needed to convince the patient to cease running. Conventional radiography rarely will be positive sooner than four weeks, by which time periosteal new bone becomes visible over stress fractures of long bones. The fracture lines are demonstrable infrequently; good quality films are essential, often several oblique views. Tomography is helpful with some fractures (for example the pars interarticularis and tarsal navicular) but CAT scans (Figure 8) provide superior diagnostic information with less irradiation; they are particularly useful in the assessment of slow healing.

Differential diagnosis

A history of stress overload, pain which is typical in site and relationship to activity and point tenderness are characteristic of stress fracture.




Figure 7. Bone scan of an early tarsal navicular stress fracture.

Differential diagnoses include osteogenic sarcoma, osteoid osteoma, osteomyelitis, osteomalacia and 'shin splints'.

'Shin splints' — a condition resulting from overuse of the tendoperiosteal tissue of the posteromedial border of the tibia — can be difficult to distinguish. Associated pain and tenderness over a large area of the tibia is exacerbated by activity and promptly relieved by rest. Radiography will be negative; a bone scan might be positive but there is increased radioactivity over 30 to 50 per cent of the tibial shaft linearly.

The two major reasons for failure to diagnose a stress fracture are neglect of it as a diagnostic possibility and acceptance of negative radiography which has been taken prematurely or examines the region inadequately.




Figure 5. Classical upper tibial stress fracture in a 13-year-old athlete.




Figure 6. Stress fracture of the left inferior pubic ramus in a 48-year-old woman jogger.




Figure 8. CAT scan of a stress fracture of the tarsal navicular bone.

Management

Removal of the cause (running) usually will result in healing within 10 weeks. Often these patients are most impatient to return to their sport and should commence other weight supported exercises such as cycling or swimming. 'Aqua-running' ('running' chest deep in a swimming pool using a floatation vest) to maintain cardiorespiratory fitness¹² may be recommended.

References

1. Breithaupt M D. Zur pathologie des mensch lichen fusses. Med Zeitung 1855; 24:169-177.
2. Stechow A W. Fussoedem und roentgenstrahlen. *Dtsch Mil-aerztl Zeitg* 1897; 26:465-471.
3. Pirker H. Die verletzungen durch muskelzug. *Ergebn Chir Orthop* 1934; 27:553-634.
4. Devas M B. Stress fractures. Edinburgh: Churchill Livingstone 1975.
5. Jackson D W, Wiltse L L, Cirincione R J. Spondylolysis in the female gymnast. *Clin Orthop* 1976; 117:68-73.
6. Whitaker J L. Arm wrestling fractures: a humerus twist. *Am J Sports Med* 1977; 5:67-77.
7. Stark H H, Jobe F W, Boyes J H, Ashworth C R. Fractures of the hook of the hamate in athletes. *J Bone Joint Surg (Am)* 1977; 59:575-582.
8. Stanitski C L, McMaster J H, Scranton P E. On the nature of stress fractures. *Am J Sports Med* 1978; 6:391-396.
9. Geslien G E, Thrall J H, Espinosa J L, Older R A. Early detection of stress fractures using 99m Tc-polyphosphate. *Radiology* 1976; 121:683-687.
10. Pavlov H, Nelson T L, Warren R F, Torg J S, Burstein A H. Stress fracture of the pubic ramus. *J Bone Joint Surg (Am)* 1982; 64:1020-1025.
11. Lombardo S J, Benson D W. Stress fractures of the femur in runners. *Am J Sports Med* 1982; 10:219-227.
12. Brody D M. Running injuries. Clinical Symposia. Summit, New Jersey: CIBA 1980; 27.

Reprint requests
Dr K D Fitch
Sports Injury Clinic
Royal Perth Rehabilitation Hospital
Shenton Park 6008


DISTRIBUTED BY HOLT-SAUNDERS PTY. LIMITED
9 Waltham Street, Artarmon 2064. (02) 439 3633

Colour Atlas of Gynaecology

Prepared by leading Australian gynaecologists,

BEISCHER & MACKAY

\$29.95



\$29.95

"For the busy G.P., this is probably the best quick reference book on gynaecological conditions in this country"

From a review in the Australian GP January/February edition 1982.

Please send me a copy of the "Colour Atlas of Gynaecology" on 30 day approval at a value of \$29.95.

NAME: _____

ADDRESS: _____

Code _____

SIGNATURE: _____ 84.131 OG01