		doc 1		doc 2	decision	id
	uvc_1				decision	Iu
			authors	Martin Meyries Roland Schnaubelt		
cases	authors	Martin Meyries Roland Schnaubelt	title	Interpolation, embeddings and traces of anisotropic fractional Sobolev spaces with temporal weights		
			publication_date   2012-02-17 00:00:00			'
	title	Interpolation, embeddings and traces of anisotropic fractional Sobolev spaces with temporal weights	source	SupportedSources.INTERNET_ARCHIVE	]	
	publication_date	2012-01-01 00:00:00	journal		]	
	source	SupportedSources.INTERNET_ARCHIVE	volume			
	journal	Elsevier BV	doi		DUPLICATES 189	
	volume		urls	https://archive.org/download/arxiv- 1202.3870/1202.3870.pdf		180
	doi	10.1016/j.jfa.2011.11.001				
		https://web.archive.org/web/20171004130229/http://publisher-				
	urls	connector.core.ac.uk/resourcesync/data/elsevier/pdf/e01/aHR0cDovL2FwaS5lbHNldmllci5jb20vY29udGVudC9hcnRpY2xlL3BpaS9zMDAyMjEyMzYxMTAwMzkyMg%3D%3D.pdf	id	id-5764273436851199179	]	
				We investigate the properties of a class of		1 1
	id	id2976003306916932408	abstract	weighted vector-valued L_p-spaces and the corresponding (an)isotropic Sobolev-Slobodetskii spaces. These spaces arise naturally in the context of maximal L_p-regularity for parabolic initial-boundary value problems. Our main tools are operators with a bounded ^â^ž-calculus, interpolation theory, and operator sums.		
	abstract	We investigate the properties of a class of weighted vector-valued L p -spaces and the corresponding (an)isotropic Sobolev-Slobodetskii spaces. These spaces arise naturally in the context of maximal L pregularity for parabolic initial-boundary value problems. Our main tools are operators with a bounded H â^žcalculus, interpolation theory, and operator sums.				
	versions					
			versions		]	