

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>BRUVERIS, MARTINS</li><li>MICHOR, PETER W.</li><li>MUMFORD, DAVID</li></ul>	authors	<ul style="list-style-type: none"><li>Martins Bruveris</li><li>Peter W. Michor</li><li>David Mumford</li></ul>	DUPLICATES	25
	title	GEODESIC COMPLETENESS FOR SOBOLEV METRICS ON THE SPACE OF IMMERSSED PLANE CURVES	title	GEODESIC COMPLETENESS FOR SOBOLEV METRICS ON THE SPACE OF IMMERSSED PLANE CURVES		
	publication_date	2017-08-02 00:00:00	publication_date	2013-12-17 00:00:00		
	source	SupportedSources.CORE	source	SupportedSources.OPENALEX		
	journal		journal	Forum of Mathematics, Sigma		
	volume		volume	2		
	doi	None	doi	10.1017/fms.2014.19		
	urls	<ul style="list-style-type: none"><li>https://core.ac.uk/download/85211737.pdf</li></ul>	urls	<ul style="list-style-type: none"><li>https://openalex.org/W2122415813</li><li>https://doi.org/10.1017/fms.2014.19</li><li>https://www.cambridge.org/core/services/aop-cambridge-core/content/view/90B8888AD48EA07AF5B02194C97D0547/S205050941400019Xa.pdf/div-class-title-geodesic-completeness-for-sobolev-metrics-on-the-space-of-immersed-plane-curves-div.pdf</li></ul>		
	id	id-5866054858517203839	id	id8021220886847111620		
	abstract	We study properties of Sobolev-type metrics on the space of immersed plane curves. We show that the geodesic equation for Sobolev-type metrics with constant coefficients of order2 and higher is globally well-posed for smooth initial data as well as for initial data in certain Sobolev spaces. Thus the space of closed plane curves equipped with such a metric is geodesically complete. We find lower bounds for the geodesic distance in terms of curvature and its derivative	abstract			
	versions		versions			