

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Evseev, N.	authors	<ul style="list-style-type: none">Nikita Evseev	NOT DUPLICATES	625
	title	Vector-valued Sobolev spaces based on Banach function spaces	title	Vector-valued Sobolev spaces based on Banach function spaces		
	publication_date	2021-01-01 00:00:00	publication_date	2020-09-21 08:50:46+00:00		
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi	10.1016/j.na.2021.112479	doi			
	urls	<ul style="list-style-type: none">https://api.elsevier.com/content/article/PII:S0362546X21001504?httpAccept=text/xmlhttps://api.elsevier.com/content/article/PII:S0362546X21001504?httpAccept=text/plainhttp://dx.doi.org/10.1016/j.na.2021.112479	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/2009.09686v1http://arxiv.org/abs/2009.09686v1http://arxiv.org/pdf/2009.09686v1		
	id	id-8875889810055077320	id	id-4899262326658041798		
	abstract		abstract	It is known that for Banach valued functions there are several approaches to define a Sobolev class. We compare the usual definition via weak derivatives with the Reshetnyak-Sobolev space and with the Newtonian space; in particular, we provide sufficient conditions when all three agree. As well we revise the difference quotient criterion and the property of Lipschitz mapping to preserve Sobolev space when it acting as a superposition operator.		
	versions		versions			