

cases	doc_1		doc_2		decision	id
					DUPLICATES	1639
	authors	<ul style="list-style-type: none">Alexander KozhevnikovAlexander G.Ramm	authors	<ul style="list-style-type: none">Alexander KozhevnikovAlexander G. Ramm		
	title	Integral Operators Basic in Random Fields Estimation Theory	title	Integral Operators Basic in Random Fields Estimation Theory		
	publication_date	2004-05-03 00:00:00	publication_date	2004-05-03 15:24:04+00:00		
	source	SupportedSources.INTERNET_ARCHIVE	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none">https://archive.org/download/arxiv-math-ph0405002/math-ph0405002.pdf	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/math-ph/0405002v1http://arxiv.org/abs/math-ph/0405002v1http://arxiv.org/pdf/math-ph/0405002v1		
	id	id-3102965136030222420	id	id1851573468644438838		
	abstract	The paper deals with the basic integral equation of random field estimation theory by the criterion of minimum of variance of the error estimate. This integral equation is of the first kind. The corresponding integra operator over a bounded domain $\hat{\mathbb{C}}$ in \mathbb{R}^n is weakly singular. This operator is an isomorphism between appropriate Sobolev spaces. This is proved by a reduction of the integral equ an elliptic boundary value problem in the domain exterior to $\hat{\mathbb{C}}$. Extra difficulties arise due to the fact that the exterior boundary value problem should be solved in the Sobolev spaces of negative order.	abstract	The paper deals with the basic integral equation of random field estimation theory by the criterion of minimum of variance of the error estimate. This integral equation is of the first kind. The corresponding integra\$ operator over a bounded domain Ω \$ in $\{\mathbb{Bbb R}\}^n$ \$ is weakly singular. This operator is an isomorphism between appropriate Sobolev spaces. This is proved by a reduction of the integral equ\$ an elliptic boundary value problem in the domain exterior to Ω .\$ Extra difficulties arise due to the fact that the exterior boundary value problem should be solved in the Sobolev spaces of negative order.		
	versions		versions			