	doc_1		doc_2		decision	id
	authors	• J. Bruna	authors	Bruna, Joaquim		
	title	L^p estimates for Riesz transforms on forms in the Poincare space H^n	title	L^p estimates for Riesz transforms on forms in the Poincare space H^n		
	publication_date	2004-11-25 00:00:00	publication_date	2004-11-25 00:00:00]	
	source	SupportedSources.SEMANTIC_SCHOLAR	source	SupportedSources.CORE	DUPLICATES 1640	
	journal	arXiv: Analysis of PDEs	journal	None		
cases	volume		volume			
	doi		doi	10.1512/iumj.2005.54.2501		
	urls	https://www.semanticscholar.org/paper/e6f1e77723da123580af3f50791715599e3908fb	urls	https://core.ac.uk/download/19577046.pdf		1640
	id	id-7641292473930828713	id	id3510497583058805521		
	abstract	Using hyperbolic form convolution with doubly isometry-invariant kernels, the explicit expression of the inverse of the de Rham laplacian acting on m-forms in the Poincar\'{e} space is found. Also, by means of some estimates for hyperbolic singular integrals, we obtain L^p -estimates for the Riesz transforms passing from the Laplacian to other covariant derivatives, in a range of p depending on m,n. Finally, using these, it is shown that the Laplacian defines topological isomorphisms in the scale of form Sobolev spaces, for m different from $n/2$, $(n+1)/2$, $(n-1)/2$.	abstract	Using hyperbolic form convolution with doubly isometry-invariant kernels, the explicit expression of the inverse of the de Rham laplacian acting on m-forms in the Poincar\'{e} space is found. Also, by means of some estimates for hyperbolic singular integrals, we obtain L^p-estimates for the Riesz transforms passing from the Laplacian to other covariant derivatives, in a range of p depending on m,n. Finally, using these, it is shown that the Laplacian defines topological isomorphisms in the scale of form Sobolev spaces, for m different from $n/2$, $(n+1)/2$, $(n-1)/2$.Comment: To appear in Indiana Univ. Math.		
	versions		versions			