

cases	doc_1		doc_2				decision	id
	authors	<ul style="list-style-type: none">Richard B. MelroseGunther Uhlmann	authors	<ul style="list-style-type: none">R. MelroseG. Uhlmann			DUPLICATES	1534
	title	Generalized backscattering and the Lax-Phillips transform	title	Generalized backscattering and the Lax-Phillips transform				
	publication_date	2007-12-01 00:00:00	publication_date	2007-12-01 00:00:00				
	source	SupportedSources.OPENALEX	source	SupportedSources.SEMANTIC_SCHOLAR				
	journal	Serdica. Mathematical Journal	journal	Serdica. Mathematical Journal				
	volume		volume					
	doi	None	doi					
	urls	<ul style="list-style-type: none">https://openalex.org/W194256293	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/3c8c7e4e26b883639a976754ffd7b390b39b0f4d				
	id	id-1813670102075724322	id	id-7138442798024459059				
	abstract		abstract	Using the free-space translation representation (modified Radon transform) of Lax and Phillips in odd dimensions, it is shown that the generalized backscattering transform (so outgoing angle $\hat{\Gamma}^{\%}_o = S\hat{\Gamma}$, in terms of the incoming angle with S orthogonal and $\text{Id}\hat{\alpha}^{\sim}S$ invertible) may be further restricted to give an entire, globally Fredholm, operator on appropriate Sobolev spaces of potentials with compact support. As a corollary we show that the modified backscattering map is a local isomorphism near elements of a generic set of potentials.				
	versions		versions					