

cases	doc_1		doc_2		decision	id
					DUPLICATES	305
	authors	<ul style="list-style-type: none">Johanna SenkKarolĀna KorvasovĀ;Jannis SchueckerEspen Hagen, Tom TetzlaffMarkus DiesmannMoritz Helias	authors	<ul style="list-style-type: none">Diesmann, MarkusHagen, EspenHelias, MoritzKorvasovĀ;, KarolĀnaSchuecker, JannisSenk, JohannaTetzlaff, Tom		
	title	Conditions for wave trains in spiking neural networks	title	Conditions for wave trains in spiking neural networks		
	publication_date	2019-09-23 00:00:00	publication_date	2019-01-01 00:00:00		
	source	SupportedSources.INTERNET_ARCHIVE	source	SupportedSources.CORE		
	journal		journal			
	volume		volume			
	doi		doi	10.1103/physrevresearch.2.023174		
	urls	<ul style="list-style-type: none">https://web.archive.org/web/20200930054246/https://arxiv.org/pdf/1801.06046v2.pdf	urls	<ul style="list-style-type: none">http://arxiv.org/abs/1801.06046		
	id	id-8141362539885483323	id	id-9157850239268272906		
	abstract	Spatiotemporal patterns such as traveling waves are frequently observed in recordings of neural activity. The mechanisms underlying the generation of such patterns are largely unknown. Previous studies have investigated the existence and uniqueness of different types of waves or bumps of activity using neural-field models, phenomenological coarse-grained descriptions of neural-network dynamics. But it remains unclear how these insights can be transferred to more biologically realistic networks of spiking neurons, where individual neurons fire irregularly. Here, we employ mean-field theory to reduce a microscopic model of leaky integrate-and-fire (LIF) neurons with distance-dependent connectivity to an effective neural-field model. In contrast to existing phenomenological descriptions, the dynamics in this neural-field model depends on the mean and the variance in the synaptic input, both determining the amplitude and the temporal structure of the resulting effective coupling kernel. For the neural-field model we employ liner stability analysis to derive conditions for the existence of spatial and temporal oscillations and wave trains, that is, temporally and spatially periodic traveling waves. We first prove that wave trains cannot occur in a single homogeneous population of neurons, irrespective of the form of distance dependence of the connection probability. Compatible with the architecture of cortical neural networks, wave trains emerge in two-population networks of excitatory and inhibitory neurons as a combination of delay-induced temporal oscillations and spatial oscillations due to distance-dependent connectivity profiles. Finally, we demonstrate quantitative agreement between predictions of the analytically tractable neural-field model and numerical simulations of both networks of nonlinear rate-based units and networks of LIF neurons.	abstract	Spatiotemporal patterns such as traveling waves are frequently observed in recordings of neural activity. The mechanisms underlying the generation of such patterns are largely unknown. Previous studies have investigated the existence and uniqueness of different types of waves or bumps of activity using neural-field models, phenomenological coarse-grained descriptions of neural-network dynamics. But it remains unclear how these insights can be transferred to more biologically realistic networks of spiking neurons, where individual neurons fire irregularly. Here, we employ mean-field theory to reduce a microscopic model of leaky integrate-and-fire (LIF) neurons with distance-dependent connectivity to an effective neural-field model. In contrast to existing phenomenological descriptions, the dynamics in this neural-field model depends on the mean and the variance in the synaptic input, both determining the amplitude and the temporal structure of the resulting effective coupling kernel. For the neural-field model we employ liner stability analysis to derive conditions for the existence of spatial and temporal oscillations and wave trains, that is, temporally and spatially periodic traveling waves. We first prove that wave trains cannot occur in a single homogeneous population of neurons, irrespective of the form of distance dependence of the connection probability. Compatible with the architecture of cortical neural networks, wave trains emerge in two-population networks of excitatory and inhibitory neurons as a combination of delay-induced temporal oscillations and spatial oscillations due to distance-dependent connectivity profiles. Finally, we demonstrate quantitative agreement between predictions of the analytically tractable neural-field model and numerical simulations of both networks of nonlinear rate-based units and networks of LIF neurons.Comment: 36 pages, 8 figures, 4 table		
	versions		versions			