

cases	doc_1		doc_2		decision	id																					
			<table><tr><td>authors</td><td><ul style="list-style-type: none">Li Xu</td></tr><tr><td>title</td><td>On the ideal magnetohydrodynamics in three-dimensional thin domains: well-posedness and asymptotics</td></tr><tr><td>publication_date</td><td>2017-07-09 08:56:21+00:00</td></tr><tr><td>source</td><td>SupportedSources.ARXIV</td></tr><tr><td>journal</td><td>None</td></tr><tr><td>volume</td><td></td></tr><tr><td>doi</td><td></td></tr><tr><td>urls</td><td><ul style="list-style-type: none">http://arxiv.org/pdf/1707.02544v2http://arxiv.org/abs/1707.02544v2http://arxiv.org/pdf/1707.02544v2</td></tr><tr><td>id</td><td>id-7479813779775126322</td></tr><tr><td>abstract</td><td>We consider the ideal magnetohydrodynamics (MHD) subjected to a strong magnetic field along x_1 direction in three-dimensional thin domains $\Omega_\delta=\mathbb{R}^2\times(-\delta,\delta)$ with slip boundary conditions. It is well-known that in this situation the system will generate Alfvén waves. Our results are summarized as follows: (i). We construct the global solutions (Alfvén waves) to MHD in the thin domain Ω_δ with $\delta>0$. In addition, the uniform energy estimates are obtained with respected to the parameter δ. (ii). We justify the asymptotics of the MHD equations from the thin domain Ω_δ to the plane \mathbb{R}^2. More precisely, we prove that the 3D Alfvén waves in Ω_δ will converge to the Alfvén waves in \mathbb{R}^2 in the limit that δ goes to zero. This shows that Alfvén waves propagating along the horizontal direction of the (3D) strip are stable and can be approximated by the (2D) Alfvén waves when δ is sufficiently small. Moreover, the control of the (2D) Alfvén waves can be obtained from the control of (3D) Alfvén waves in the thin domain Ω_δ with aid of the uniform bounds. The proofs of main results rely on the design of the proper energy functional and the null structures of the nonlinear terms. Here the null structures means two aspects: separation of the Alfvén waves (z_+ and z_-) and no bad quadratic terms $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ where $z_\pm=(z\pm z^3_\pm)^h$ and $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ is the linear combination of terms $\partial^\alpha\partial_{z_-}^h\partial^\beta\partial_{z_+}^h$ with $\alpha, \beta\in (\mathbb{Z}_-\geq 0)^3$.</td></tr><tr><td>versions</td><td></td></tr></table>	authors	<ul style="list-style-type: none">Li Xu	title	On the ideal magnetohydrodynamics in three-dimensional thin domains: well-posedness and asymptotics	publication_date	2017-07-09 08:56:21+00:00	source	SupportedSources.ARXIV	journal	None	volume		doi		urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1707.02544v2http://arxiv.org/abs/1707.02544v2http://arxiv.org/pdf/1707.02544v2	id	id-7479813779775126322	abstract	We consider the ideal magnetohydrodynamics (MHD) subjected to a strong magnetic field along x_1 direction in three-dimensional thin domains $\Omega_\delta=\mathbb{R}^2\times(-\delta,\delta)$ with slip boundary conditions. It is well-known that in this situation the system will generate Alfvén waves. Our results are summarized as follows: (i). We construct the global solutions (Alfvén waves) to MHD in the thin domain Ω_δ with $\delta>0$. In addition, the uniform energy estimates are obtained with respected to the parameter δ . (ii). We justify the asymptotics of the MHD equations from the thin domain Ω_δ to the plane \mathbb{R}^2 . More precisely, we prove that the 3D Alfvén waves in Ω_δ will converge to the Alfvén waves in \mathbb{R}^2 in the limit that δ goes to zero. This shows that Alfvén waves propagating along the horizontal direction of the (3D) strip are stable and can be approximated by the (2D) Alfvén waves when δ is sufficiently small. Moreover, the control of the (2D) Alfvén waves can be obtained from the control of (3D) Alfvén waves in the thin domain Ω_δ with aid of the uniform bounds. The proofs of main results rely on the design of the proper energy functional and the null structures of the nonlinear terms. Here the null structures means two aspects: separation of the Alfvén waves (z_+ and z_-) and no bad quadratic terms $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ where $z_\pm=(z\pm z^3_\pm)^h$ and $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ is the linear combination of terms $\partial^\alpha\partial_{z_-}^h\partial^\beta\partial_{z_+}^h$ with $\alpha, \beta\in (\mathbb{Z}_-\geq 0)^3$.	versions		DUPLICATES	128
	authors	<ul style="list-style-type: none">Li Xu																									
	title	On the ideal magnetohydrodynamics in three-dimensional thin domains: well-posedness and asymptotics																									
	publication_date	2017-07-09 08:56:21+00:00																									
	source	SupportedSources.ARXIV																									
	journal	None																									
	volume																										
	doi																										
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1707.02544v2http://arxiv.org/abs/1707.02544v2http://arxiv.org/pdf/1707.02544v2																									
	id	id-7479813779775126322																									
	abstract	We consider the ideal magnetohydrodynamics (MHD) subjected to a strong magnetic field along x_1 direction in three-dimensional thin domains $\Omega_\delta=\mathbb{R}^2\times(-\delta,\delta)$ with slip boundary conditions. It is well-known that in this situation the system will generate Alfvén waves. Our results are summarized as follows: (i). We construct the global solutions (Alfvén waves) to MHD in the thin domain Ω_δ with $\delta>0$. In addition, the uniform energy estimates are obtained with respected to the parameter δ . (ii). We justify the asymptotics of the MHD equations from the thin domain Ω_δ to the plane \mathbb{R}^2 . More precisely, we prove that the 3D Alfvén waves in Ω_δ will converge to the Alfvén waves in \mathbb{R}^2 in the limit that δ goes to zero. This shows that Alfvén waves propagating along the horizontal direction of the (3D) strip are stable and can be approximated by the (2D) Alfvén waves when δ is sufficiently small. Moreover, the control of the (2D) Alfvén waves can be obtained from the control of (3D) Alfvén waves in the thin domain Ω_δ with aid of the uniform bounds. The proofs of main results rely on the design of the proper energy functional and the null structures of the nonlinear terms. Here the null structures means two aspects: separation of the Alfvén waves (z_+ and z_-) and no bad quadratic terms $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ where $z_\pm=(z\pm z^3_\pm)^h$ and $Q(\partial_{z_-}^h, \partial_{z_+}^h)$ is the linear combination of terms $\partial^\alpha\partial_{z_-}^h\partial^\beta\partial_{z_+}^h$ with $\alpha, \beta\in (\mathbb{Z}_-\geq 0)^3$.																									
	versions																										
	<table><tr><td>authors</td><td><ul style="list-style-type: none">Li Xu</td></tr><tr><td>title</td><td>On the Ideal Magnetohydrodynamics in Three-Dimensional Thin Domains: Well-Posedness and Asymptotics</td></tr><tr><td>publication_date</td><td>2017-07-09 00:00:00</td></tr><tr><td>source</td><td>SupportedSources.SEMANTIC_SCHOLAR</td></tr><tr><td>journal</td><td>Archive for Rational Mechanics and Analysis</td></tr><tr><td>volume</td><td>236</td></tr><tr><td>doi</td><td>10.1007/s00205-019-01464-8</td></tr><tr><td>urls</td><td><ul style="list-style-type: none">https://www.semanticscholar.org/paper/bc74b65c00c87bcf38161269df6ebe1e0c510398</td></tr><tr><td>id</td><td>id1995204750088286288</td></tr><tr><td>abstract</td><td>None</td></tr><tr><td>versions</td><td></td></tr></table>	authors	<ul style="list-style-type: none">Li Xu	title	On the Ideal Magnetohydrodynamics in Three-Dimensional Thin Domains: Well-Posedness and Asymptotics	publication_date	2017-07-09 00:00:00	source	SupportedSources.SEMANTIC_SCHOLAR	journal	Archive for Rational Mechanics and Analysis	volume	236	doi	10.1007/s00205-019-01464-8	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/bc74b65c00c87bcf38161269df6ebe1e0c510398	id	id1995204750088286288	abstract	None	versions					
	authors	<ul style="list-style-type: none">Li Xu																									
	title	On the Ideal Magnetohydrodynamics in Three-Dimensional Thin Domains: Well-Posedness and Asymptotics																									
	publication_date	2017-07-09 00:00:00																									
	source	SupportedSources.SEMANTIC_SCHOLAR																									
	journal	Archive for Rational Mechanics and Analysis																									
	volume	236																									
doi	10.1007/s00205-019-01464-8																										
urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/bc74b65c00c87bcf38161269df6ebe1e0c510398																										
id	id1995204750088286288																										
abstract	None																										
versions																											