

cases	doc_1		doc_2		decision	id
			authors	<ul style="list-style-type: none">Li Xu	NOT DUPLICATES	1007
	title	On the Ideal Magnetohydrodynamics in Three-Dimensional Thin Domains: Well-Posedness and Asymptotics	title	On the ideal magnetohydrodynamics in three-dimensional thin domains: well-posedness and asymptotics		
	publication_date	2019-11-08 00:00:00	publication_date	2017-07-09 08:56:21+00:00		
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi	10.1007/s00205-019-01464-8	doi			
	urls	<ul style="list-style-type: none">http://link.springer.com/content/pdf/10.1007/s00205-019-01464-8.pdfhttp://link.springer.com/article/10.1007/s00205-019-01464-8/fulltext.htmlhttp://link.springer.com/content/pdf/10.1007/s00205-019-01464-8.pdfhttp://dx.doi.org/10.1007/s00205-019-01464-8	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1707.02544v2http://arxiv.org/abs/1707.02544v2http://arxiv.org/pdf/1707.02544v2		
	id	id-3953186404005294396	id	id-7479813779775126322		
	abstract		abstract	We consider the ideal magnetohydrodynamics (MHD) subjected to a strong magnetic field along x_1 direction in three-dimensional thin domains $\Omega_\delta=\mathbb{R}^2\times(-\delta,\delta)$ with slip boundary conditions. It is well-known that in this situation the system will generate Alfvén waves. Our results are summarized as follows: (i)., We construct the global solutions (Alfvén waves) to MHD in the thin domain Ω_δ with $\delta>0$. In addition, the uniform energy estimates are obtained with respected to the parameter δ . (ii). We justify the asymptotics of the MHD equations from the thin domain Ω_δ to the plane \mathbb{R}^2 . More precisely, we prove that the 3D Alfvén waves in Ω_δ will converge to the Alfvén waves in \mathbb{R}^2 in the limit that δ goes to zero. This shows that Alfvén waves propagating along the horizontal direction of the (3D) strip are stable and can be approximated by the (2D) Alfvén waves when δ is sufficiently small. Moreover, the control of the (2D) Alfvén waves can be obtained from the control of (3D) Alfvén waves in the thin domain Ω_δ with aid of the uniform bounds. The proofs of main results rely on the design of the proper energy functional and the null structures of the nonlinear terms. Here the null structures means two aspects: separation of the Alfvén waves (z_+ and z_-) and no bad quadratic terms $Q(\partial_3 z_-^h, \partial_3 z_+^h)$ where $z_\pm=(z_\pm^h, z_\pm^3)$ and $Q(\partial_3 z_-^h, \partial_3 z_+^h)$ is the linear combination of terms $\partial^\alpha \partial_3 z_-^h \partial^\beta \partial_3 z_+^h$ with $\alpha, \beta \in (\mathbb{Z})_{\geq 0}^3$.		
	versions		versions			