

cases	doc_1		doc_2		decision	id																																											
	<div></div> <table><tr><td>authors</td><td><ul style="list-style-type: none">J. Thim</td></tr><tr><td>title</td><td>Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant</td></tr><tr><td>publication_date</td><td>2015-01-23 00:00:00</td></tr><tr><td>source</td><td>SupportedSources.SEMANTIC_SCHOLAR</td></tr><tr><td>journal</td><td>Potential Analysis</td></tr><tr><td>volume</td><td>43</td></tr><tr><td>doi</td><td>10.1007/s11118-015-9464-7</td></tr><tr><td>urls</td><td><ul style="list-style-type: none">https://www.semanticscholar.org/paper/dea53a46a82077fc7e2c62dcc0c789128fc3258d</td></tr><tr><td>id</td><td>id4721265325280768</td></tr><tr><td>abstract</td><td>None</td></tr><tr><td>versions</td><td></td></tr></table>		authors	<ul style="list-style-type: none">J. Thim	title	Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant	publication_date	2015-01-23 00:00:00	source	SupportedSources.SEMANTIC_SCHOLAR	journal	Potential Analysis	volume	43	doi	10.1007/s11118-015-9464-7	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/dea53a46a82077fc7e2c62dcc0c789128fc3258d	id	id4721265325280768	abstract	None	versions		<table><tr><td>authors</td><td><ul style="list-style-type: none">Johan Thim</td></tr><tr><td>title</td><td>Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant</td></tr><tr><td>publication_date</td><td>2014-05-08 23:13:15+00:00</td></tr><tr><td>source</td><td>SupportedSources.ARXIV</td></tr><tr><td>journal</td><td>None</td></tr><tr><td>volume</td><td></td></tr><tr><td>doi</td><td></td></tr><tr><td>urls</td><td><ul style="list-style-type: none">http://arxiv.org/pdf/1405.2121v1http://arxiv.org/abs/1405.2121v1http://arxiv.org/pdf/1405.2121v1</td></tr><tr><td>id</td><td>id6216282122430386478</td></tr><tr><td>abstract</td><td>This article considers two weight estimates for the single layer potential --- corresponding to the Laplace operator in \mathbb{R}^{N+1} --- on Lipschitz surfaces with small Lipschitz constant. We present conditions on the weights to obtain solvability and uniqueness results in weighted Lebesgue spaces and weighted homogeneous Sobolev spaces, where the weights are assumed to be radial and doubling. In the case when the weights are additionally assumed to be differentiable almost everywhere, simplified conditions in terms of the logarithmic derivative are presented, and as an application, we prove that the operator corresponding to the single layer potential in question is an isomorphism between certain weighted spaces of the type mentioned above. Furthermore, we consider several explicit weight functions. In particular, we present results for power exponential weights which generalize known results for the case when the single layer potential is reduced to a Riesz potential, which is the case when the Lipschitz surface is given by a hyperplane.</td></tr><tr><td>versions</td><td></td></tr></table>	authors	<ul style="list-style-type: none">Johan Thim	title	Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant	publication_date	2014-05-08 23:13:15+00:00	source	SupportedSources.ARXIV	journal	None	volume		doi		urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1405.2121v1http://arxiv.org/abs/1405.2121v1http://arxiv.org/pdf/1405.2121v1	id	id6216282122430386478	abstract	This article considers two weight estimates for the single layer potential --- corresponding to the Laplace operator in \mathbb{R}^{N+1} --- on Lipschitz surfaces with small Lipschitz constant. We present conditions on the weights to obtain solvability and uniqueness results in weighted Lebesgue spaces and weighted homogeneous Sobolev spaces, where the weights are assumed to be radial and doubling. In the case when the weights are additionally assumed to be differentiable almost everywhere, simplified conditions in terms of the logarithmic derivative are presented, and as an application, we prove that the operator corresponding to the single layer potential in question is an isomorphism between certain weighted spaces of the type mentioned above. Furthermore, we consider several explicit weight functions. In particular, we present results for power exponential weights which generalize known results for the case when the single layer potential is reduced to a Riesz potential, which is the case when the Lipschitz surface is given by a hyperplane.	versions		DUPLICATES	1113
			authors	<ul style="list-style-type: none">J. Thim																																													
			title	Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant																																													
			publication_date	2015-01-23 00:00:00																																													
			source	SupportedSources.SEMANTIC_SCHOLAR																																													
			journal	Potential Analysis																																													
			volume	43																																													
			doi	10.1007/s11118-015-9464-7																																													
			urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/dea53a46a82077fc7e2c62dcc0c789128fc3258d																																													
			id	id4721265325280768																																													
	abstract	None																																															
	versions																																																
	authors	<ul style="list-style-type: none">Johan Thim																																															
	title	Two Weight Estimates for the Single Layer Potential on Lipschitz Surfaces with Small Lipschitz Constant																																															
	publication_date	2014-05-08 23:13:15+00:00																																															
	source	SupportedSources.ARXIV																																															
	journal	None																																															
	volume																																																
	doi																																																
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1405.2121v1http://arxiv.org/abs/1405.2121v1http://arxiv.org/pdf/1405.2121v1																																															
id	id6216282122430386478																																																
abstract	This article considers two weight estimates for the single layer potential --- corresponding to the Laplace operator in \mathbb{R}^{N+1} --- on Lipschitz surfaces with small Lipschitz constant. We present conditions on the weights to obtain solvability and uniqueness results in weighted Lebesgue spaces and weighted homogeneous Sobolev spaces, where the weights are assumed to be radial and doubling. In the case when the weights are additionally assumed to be differentiable almost everywhere, simplified conditions in terms of the logarithmic derivative are presented, and as an application, we prove that the operator corresponding to the single layer potential in question is an isomorphism between certain weighted spaces of the type mentioned above. Furthermore, we consider several explicit weight functions. In particular, we present results for power exponential weights which generalize known results for the case when the single layer potential is reduced to a Riesz potential, which is the case when the Lipschitz surface is given by a hyperplane.																																																
versions																																																	