

cases	doc_1		doc_2		decision	id
					DUPLICATES	120
	authors	<ul style="list-style-type: none">Zoltan MuzsnayPeter T. Nagy	authors	<ul style="list-style-type: none">Zoltan MuzsnayPeter T. Nagy		
	title	Finsler spaces with infinite dimensional holonomy group	title	Finsler spaces with infinite dimensional holonomy group		
	publication_date	2010-12-02 00:00:00	publication_date	2010-12-02 19:31:20+00:00		
	source	SupportedSources.INTERNET_ARCHIVE	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none">https://archive.org/download/arxiv-1012.0545/1012.0545.pdf	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1012.0545v1http://arxiv.org/abs/1012.0545v1http://arxiv.org/pdf/1012.0545v1		
	id	id7118543011671501685	id	id886960732783481477		
	abstract	Our paper is devoted to the study of the holonomy groups of Finsler surfaces using the methods of infinite dimensional Lie theory. The notion of infinitesimal holonomy algebra will be introduced, by the smallest Lie algebra of vector fields on an indicatrix, containing the curvature vector fields and their horizontal covariant derivatives with respect to the Berwald connection. We obtain that the topological closure of the holonomy group contains the exponential image of any tangent Lie algebra of the holonomy group. A class of Randers surfaces is determined, for which the infinitesimal holonomy algebra coincides with the curvature algebra. We prove that for all projectively flat Randers surfaces of non-zero constant flag curvature the infinitesimal holonomy algebra has infinite dimension and hence the holonomy group cannot be a Lie group of finite dimension. Finally, in the case of the Funk metric we prove that the infinitesimal holonomy algebra is a dense subalgebra of the Lie algebra of the full diffeomorphism group and hence the topological closure of the holonomy group is the orientation preserving diffeomorphism group of the circle.	abstract	Our paper is devoted to the study of the holonomy groups of Finsler surfaces using the methods of infinite dimensional Lie theory. The notion of infinitesimal holonomy algebra will be introduced, by the smallest Lie algebra of vector fields on an indicatrix, containing the curvature vector fields and their horizontal covariant derivatives with respect to the Berwald connection. We obtain that the topological closure of the holonomy group contains the exponential image of any tangent Lie algebra of the holonomy group. A class of Randers surfaces is determined, for which the infinitesimal holonomy algebra coincides with the curvature algebra. We prove that for all projectively flat Randers surfaces of non-zero constant flag curvature the infinitesimal holonomy algebra has infinite dimension and hence the holonomy group cannot be a Lie group of finite dimension. Finally, in the case of the Funk metric we prove that the infinitesimal holonomy algebra is a dense subalgebra of the Lie algebra of the full diffeomorphism group and hence the topological closure of the holonomy group is the orientation preserving diffeomorphism group of the circle.		
	versions		versions			