

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Nguyen Huu Huy PhucMonica Torres	authors	<ul style="list-style-type: none">N. PhucM. Torres	DUPLICATES	1089
	title	Characterizations of signed measures in the dual of BV and related isometric isomorphisms	title	Characterizations of signed measures in the dual of BV and related isometric isomorphisms		
	publication_date	2017-03-09 00:00:00	publication_date	2015-03-20 00:00:00		
	source	SupportedSources.OPENALEX	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	Annali della Scuola normale superiore di Pisa. Classe di scienze	journal	arXiv: Analysis of PDEs		
	volume		volume			
	doi	10.2422/2036-2145.201508_005	doi			
	urls	<ul style="list-style-type: none">https://openalex.org/W2963766253https://doi.org/10.2422/2036-2145.201508_005http://arxiv.org/pdf/1503.06208	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/810b49825e3b5bcb76bcd34adc07d5cc9b2bd125		
	id	id1277234806570840108	id	id711896265102562653		
	abstract		abstract	We characterize all (signed) measures in $BV_{\frac{n}{n-1}}(\mathbb{R}^n)^*$, where $BV_{\frac{n}{n-1}}(\mathbb{R}^n)$ is defined as the space of all functions u in $L^{\frac{n}{n-1}}(\mathbb{R}^n)$ such that Du is a finite vector-valued measure. We also show that $BV_{\frac{n}{n-1}}(\mathbb{R}^n)^*$ and $BV(\mathbb{R}^n)^*$ are isometrically isomorphic, where $BV(\mathbb{R}^n)$ is defined as the space of all functions u in $L^1(\mathbb{R}^n)$ such that Du is a finite vector-valued measure. As a consequence of our characterizations, an old issue raised in Meyers-Ziemer [MZ] is resolved by constructing a locally integrable function f such that f belongs to $BV(\mathbb{R}^n)^*$ but $ f $ does not. Moreover, we show that the measures in $BV_{\frac{n}{n-1}}(\mathbb{R}^n)^*$ coincide with the measures in $\dot{W}^{1,1}(\mathbb{R}^n)^*$, the dual of the homogeneous Sobolev space $\dot{W}^{1,1}(\mathbb{R}^n)$, in the sense of isometric isomorphism. For a bounded open set Ω with Lipschitz boundary, we characterize the measures in the dual space $BV_0(\Omega)^*$. One of the goals of this paper is to make precise the definition of $BV_0(\Omega)$, which is the space of functions of bounded variation with zero trace on the boundary of Ω . We show that the measures in $BV_0(\Omega)^*$ coincide with the measures in $W^{1,1}_0(\Omega)^*$. Finally, the class of finite measures in $BV(\Omega)^*$ is also characterized.		
	versions		versions			