	doc_1		doc_2		decision
			authors	Dimitri Cobb Francesco Fanelli	
cases		• D. Cobb		Symmetry breaking in ideal magnetohydrodynamics: the role of the velocity	
	authors	• F. Fanelli	publication_date 2021-02-26 16:51:37+00:00		
			source	SupportedSources.ARXIV	
	title	Symmetry breaking in ideal magnetohydrodynamics: the role of the velocity	journal	None	
	publication_date 2021-02-26 00:00:00		volume		il I
	source	SupportedSources.SEMANTIC_SCHOLAR	doi		DUPLICATES 967
	journal	Journal of Elliptic and Parabolic Equations	urls	• http://arxiv.org/aos/2102.13360v1	
	volume	7			
	doi	10.1007/s41808-021-00105-0		• http://arxiv.org/pdf/2102.13586v1	
	urls	https://www.semanticscholar.org/paper/178932f4be021ccebd912929e77154e29e2581dc	id	id-43925019468905220	
	id	id1262168820855765753	abstract	The ideal magnetohydrodynamic equations are, roughly speaking, a quasi-linear symmetric hyperbolic system of PDEs, but not all the unknowns play the same role in this system. Indeed, in the regime of small magnetic fields, the equations are close to the incompressible Euler equations. In the present paper, we adopt this point of view to study questions linked with the lifespan of strong solutions to the ideal magnetohydrodynamic equations. First of all, we prove a continuation criterion in terms of the velocity field only. Secondly, we refine the explicit lower bound for the lifespan of \$2\$-D flows found in [11], by relaxing the regularity assumptions on the initial magnetic field.	
	abstract	None			
	versions				
			versions		