

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Roberto CastelliJean-Philippe Lessard	authors	<ul style="list-style-type: none">R. CastelliJ. Lessard	DUPLICATES	402
	title	Rigorous numerics in Floquet theory: computing stable and unstable bundles of periodic orbits	title	Rigorous Numerics in Floquet Theory: Computing Stable and Unstable Bundles of Periodic Orbits		
	publication_date	2011-12-20 23:00:29+00:00	publication_date	2011-12-20 00:00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	None	journal	SIAM J. Appl. Dyn. Syst.		
	volume		volume	12		
	doi		doi	10.1137/120873960		
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1112.4874v1http://arxiv.org/abs/1112.4874v1http://arxiv.org/pdf/1112.4874v1	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/c3e5cc8551109a6a75b6c4301b96fa71776a52a1		
	id	id1264179137467551809	id	id-927241914778945999		
	abstract	In this paper, a new rigorous numerical method to compute fundamental matrix solutions of non-autonomous linear differential equations with periodic coefficients is introduced. Decomposing the fundamental matrix solutions $\Phi(t)$ by their Floquet normal forms, that is as product of real periodic and exponential matrices $\Phi(t)=Q(t)e^{\{Rt\}}$, one solves simultaneously for R and for the Fourier coefficients of Q via a fixed point argument in a suitable Banach space of rapidly decaying coefficients. As an application, the method is used to compute rigorously stable and unstable bundles of periodic orbits of vector fields. Examples are given in the context of the Lorenz equations and the ζ^3 -model.	abstract	In this paper, a rigorous method to compute Floquet normal forms of fundamental matrix solutions of nonautonomous linear differential equations with periodic coefficients is introduced. The Floquet normal form of a fundamental matrix solution $\Phi(t)$ is a canonical decomposition of the form $\Phi(t)=Q(t)e^{\{Rt\}}$, where $Q(t)$ is a real periodic matrix and R is a constant matrix. To rigorously compute the Floquet normal form, the idea is to use the regularity of $Q(t)$ and to simultaneously solve for R and $Q(t)$ with the contraction mapping theorem in a Banach space of rapidly decaying coefficients. The explicit knowledge of R and Q can then be used to construct, in a rigorous computer-assisted way, stable and unstable bundles of periodic orbits of vector fields. The new proposed method does not require rigorous numerical integration of the ODE.		
	versions		versions			