

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Anna AnopTetiana Kasirenko	authors	<ul style="list-style-type: none">A. AnopT. Kasirenko	DUPLICATES	1110
	title	Elliptic boundary-value problems in H^{∞} -rmander spaces	title	Elliptic boundary-value problems in H^{∞} -rmander spaces		
	publication_date	2016-12-13 14:56:15+00:00	publication_date	2016-12-13 00:00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	Methods Funct. Anal. Topology, Vol. 22 (2016), no. 4, 295-310	journal			
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1612.04220v2http://arxiv.org/abs/1612.04220v2http://arxiv.org/pdf/1612.04220v2	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/653ba00f8d7b5ecd6f37d133834b113259ca1cde		
	id	id4031004533228549517	id	id-5885306826384845294		
	abstract	We investigate general elliptic boundary-value problems in H^{∞} -ormander inner product spaces that form the extended Sobolev scale. The latter consists of all Hilbert spaces that are interpolation spaces with respect to the Sobolev Hilbert scale. We prove that the operator corresponding to an arbitrary elliptic problem is Fredholm in appropriate couples of the H^{∞} -ormander spaces and induces a collection of isomorphisms on the extended Sobolev scale. We obtain a local a priory estimate for generalized solutions to this problem and prove a theorem on their local regularity in the H^{∞} -ormander spaces. We find new sufficient conditions under which generalized derivatives (of a given order) of the solutions are continuous.	abstract	We investigate general elliptic boundary-value problems in H^{∞} -rmander inner product spaces that form the extended Sobolev scale. The latter consists of all Hilbert spaces that are interpolation spaces with respect to the Sobolev Hilbert scale. We prove that the operator corresponding to an arbitrary elliptic problem is Fredholm in appropriate couples of the H^{∞} -rmander spaces and induces a collection of isomorphisms on the extended Sobolev scale. We obtain a local a priory estimate for generalized solutions to this problem and prove a theorem on their local regularity in the H^{∞} -rmander spaces. We find new sufficient conditions under which generalized derivatives (of a given order) of the solutions are continuous.		
	versions		versions			