

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Hamid SaleemZain H. Saleem	authors	<ul style="list-style-type: none">Hamid SaleemZain H. Saleem	NOT DUPLICATES	1217
	title	3-D exact analytical solutions of two fluid plasma, MHD and neutral fluid equations for the creation of ordered structures as well as jet-like flows	title	3D Exact Analytical Solutions of Two-fluid Plasma, Magnetohydrodynamics, and Neutral Fluid Equations for the Creation of Ordered Structures as well as Jet-like Flows		
	publication_date	2022-01-22 00:00:00	publication_date	2022-03-01 00:00:00		
	source	SupportedSources.INTERNET_ARCHIVE	source	SupportedSources.INTERNET_ARCHIVE		
	journal		journal	American Astronomical Society		
	volume		volume			
	doi		doi	10.3847/1538-4357/ac4d8d		
	urls	<ul style="list-style-type: none">https://web.archive.org/web/20220128052230/https://arxiv.org/pdf/2111.03927v3.pdf	urls	<ul style="list-style-type: none">https://web.archive.org/web/20220516025714/https://iopscience.iop.org/article/10.3847/1538-4357/ac4d8d/pdf		
	id	id4316993057341677641	id	id-3738786662911198726		
	abstract	The 3-D exact analytical solutions of ideal two fluid plasma, single fluid plasma (MHD) and neutral fluid equations have been found using physically justifiable assumptions. Surprisingly these solutions satisfy all non-linearities in the systems. It is pointed out that these solutions explain the fundamental mechanism behind the creation of vast variety of ordered structures in plasmas and fluids. In the limiting case of two dimensional (2-D) dependence of fields, the theoretical model for plasma is applied to explain the formation of spicules in solar chromosphere. It is pointed out that the main contribution of electron (ion) baro clinic vectors is to produce vorticity in the plasma and that magnetic field generation is coupled with the flow of both electrons and ions.	abstract	The 3D exact analytical solutions of ideal two-fluid plasma, single-fluid plasma, and neutral fluid equations have been found using physically justifiable assumptions. Surprisingly these solutions satisfy all nonlinearities in the systems. It is pointed out that these solutions explain the fundamental mechanism behind the creation of a vast variety of ordered structures in plasmas and fluids. In the limiting case of 2D dependence of fields, the theoretical model for plasma is applied to explain the formation of spicules in the solar chromosphere. It is pointed out that the main contribution of electron (ion) baroclinic vectors is to produce vorticity in the plasma, and that magnetic field generation is coupled with the flow of both electrons and ions.		
	versions		versions			