

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Karak, N.	authors	<ul style="list-style-type: none">Nijjwal Karak	NOT DUPLICATES	1186
	title	Removable Sets for Orlicz-Sobolev Spaces	title	Removable sets for Orlicz-Sobolev spaces		
	publication_date	2015-07-07 00:00:00	publication_date	2014-08-25 10:44:14+00:00		
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi	10.1007/s11118-015-9491-4	doi			
	urls	<ul style="list-style-type: none">http://link.springer.com/content/pdf/10.1007/s11118-015-9491-4.pdfhttp://link.springer.com/article/10.1007/s11118-015-9491-4/fulltext.htmlhttp://link.springer.com/content/pdf/10.1007/s11118-015-9491-4http://dx.doi.org/10.1007/s11118-015-9491-4	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1408.5716v1http://arxiv.org/abs/1408.5716v1http://arxiv.org/pdf/1408.5716v1		
	id	id-5907151826585886922	id	id-6358241440876038377		
	abstract		abstract	We study removable sets for the Orlicz-Sobolev space $W^{1,\Psi}$, for functions of the form $\Psi(t)=t^p\log^{\lambda}(e+t)$. We show that (p,λ) -porous sets lying in a hyperplane are removable and that this result is essentially sharp.		
	versions		versions			