title Dissipation_date 2001-1  source Support journal volume doi  urls  id id4901	R. Turkmani     U. Torkelsson (Chalmers University of Technology/Goteborg University)  sipation of non-linear circularly polarized Alfven waves  01-11-05 00:00:00  pportedSources.INTERNET_ARCHIVE	authors  title publication_dat source journal volume doi	R. Turkmani     U. Torkelsson  Dissipation of non-linear circularly polarized Alfven waves  te 2001-11-05 09:27:52+00:00  SupportedSources.ARXIV  None	
publication_date   2001-1   source   Support   journal     volume     doi     urls   id   id4901	01-11-05 00:00:00	publication_dat source journal volume	te 2001-11-05 09:27:52+00:00 SupportedSources.ARXIV	
publication_date   2001-1   source   Support   journal     volume     doi     urls   id   id4901	01-11-05 00:00:00	source journal volume	SupportedSources.ARXIV	
source Suppor journal volume doi urls id id4901		journal volume	^^	
journal volume doi urls id id4901	oportedSources.INTERNET_ARCHIVE	volume	None State of the	
volume doi urls id id4901				
doi urls id id4901		doi		
urls   • 1     id4901				.m. II
id id4901			http://arxiv.org/pdf/astro-ph/0111073v1	DUPLICATES 587
	<ul> <li>https://archive.org/download/arxiv-astro-ph0111073/astro-ph0111073.pdf</li> </ul>	urls	<ul> <li>http://arxiv.org/abs/astro-ph/0111073v1</li> <li>http://arxiv.org/pdf/astro-ph/0111073v1</li> </ul>	
	901243783053391397			
	We study propagating Alfven waves by solving the time-dependent equations of magnetohydrodynamics (MHD) in one dimension numerically. In a homogeneous medium the circularly polarized Alfven wave is an exact solution of the ideal MHD equations, and therefore it does not suffer from any dissipation. A high-amplitude linearly polarized Alfven wave, on the other hand, steepens and form current sheets, in which the Poynting flux is lost. In a stratified medium, however, a high-amplitude circularly polarized Alfven wave can also lose a significant fraction of its Poynting flux.	id	id6870159745096156641	
abstract exact s ampliti Poynti also lo		abstract	We study propagating Alfven waves by solving the time-dependent equations of magnetohydrodynamics (MHD) in one dimension numerically. In a homogeneous medium the circularly polarized Alfven wave is an exact solution of the ideal MHD equations, and therefore it does not suffer from any dissipation. A high-amplitude linearly polarized Alfven wave, on the other hand, steepens and form current sheets, in which the Poynting flux is lost. In a stratified medium, however, a high-amplitude circularly polarized Alfven wave can also lose a significant fraction of its Poynting flux.	
versions		versions	also lose a significant fraction of its Poynting flux.	