

cases	doc_1		doc_2		decision	id
					DUPLICATES	1118
	authors	<ul style="list-style-type: none">Downes, TurloughO\u27Sullivan, Stephen	authors	<ul style="list-style-type: none">Stephen O'SullivanTurlough P. Downes		
	title	An Explicit Scheme for Multifluid Magnetohydrodynamics	title	An explicit scheme for multifluid magnetohydrodynamics		
	publication_date	2006-01-01 00:00:00	publication_date	2005-11-16 14:18:44+00:00		
	source	SupportedSources.CORE	source	SupportedSources.ARXIV		
	journal		journal	Mon.Not.Roy.Astron.Soc.366:1329-1336,2006		
	volume		volume			
	doi	None	doi	10.1111/j.1365-2966.2005.09898.x		
	urls	<ul style="list-style-type: none">https://core.ac.uk/download/301312394.pdf	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/astro-ph/0511478v3http://dx.doi.org/10.1111/j.1365-2966.2005.09898.xhttp://arxiv.org/abs/astro-ph/0511478v3http://arxiv.org/pdf/astro-ph/0511478v3		
	id	id-4372164028258632477	id	id-3793703646682670713		
	abstract	When modelling astrophysical fluid flows, it is often appropriate to discard the canonical magnetohydrodynamic approximation, thereby freeing the magnetic field to diffuse with respect to the bulk velocity field. As a consequence, however, the induction equation can become problematic to solve via standard explicit techniques. In particular, the Hall diffusion term admits fast-moving whistler waves which can impose a vanishing time-step limit.Within an explicit differencing framework, a multifluid scheme for weakly ionized plasmas is presented which relies upon a new approach to integrating the induction equation efficiently. The first component of this approach is a relatively unknown method of accelerating the integration of parabolic systems by enforcing stability over large compound time-steps rather than over each of the constituent substeps. This method, Super Time-Stepping, proves to be very effective in applying a part of the Hall term up to a known critical value. The excess of the Hall term above this critical value is then included via a new scheme for pure Hall diffusion	abstract	When modeling astrophysical fluid flows, it is often appropriate to discard the canonical magnetohydrodynamic approximation thereby freeing the magnetic field to diffuse with respect to the bulk velocity field. As a consequence, however, the induction equation can become problematic to solve via standard explicit techniques. In particular, the Hall diffusion term admits fast-moving whistler waves which can impose a vanishing timestep limit. Within an explicit differencing framework, a multifluid scheme for weakly ionised plasmas is presented which relies upon a new approach to integrating the induction equation efficiently. The first component of this approach is a relatively unknown method of accelerating the integration of parabolic systems by enforcing stability over large compound timesteps rather than over each of the constituent substeps. This method, Super Time Stepping, proves to be very effective in applying a part of the Hall term up to a known critical value. The excess of the Hall term above this critical value is then included via a new scheme for pure Hall diffusion.		
	versions		versions			