

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Maximiliano AltamiranoRoberto CortezMatthieu JonckheereLasse Leskelä	authors	<ul style="list-style-type: none">Maximiliano AltamiranoRoberto CortezMatthieu JonckheereLasse Leskelä	DUPLICATES	47
	title	Persistence in a large network of locally interacting neurons	title	Persistence in a large network of locally interacting neurons		
	publication_date	2021-08-13 20:26:26+00:00	publication_date	2021-08-13 00:00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.INTERNET_ARCHIVE		
	journal	None	journal			
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/2108.06386v1http://arxiv.org/abs/2108.06386v1http://arxiv.org/pdf/2108.06386v1	urls	<ul style="list-style-type: none">https://web.archive.org/web/20210820005731/https://arxiv.org/pdf/2108.06386v1.pdf		
	id	id3991795243743126557	id	id-2460419395411124320		
	abstract	This article presents a biological neural network model driven by inhomogeneous Poisson processes accounting for the intrinsic randomness of synapses. The main novelty is the introduction of local interactions: each firing neuron triggers an instantaneous increase in electric potential to a fixed number of randomly chosen neurons. We prove that, as the number of neurons approaches infinity, the finite network converges to a nonlinear meanfield process characterised by a jump-type stochastic differential equation. We show that this process displays a phase transition: the activity of a typical neuron in the infinite network either rapidly dies out, or persists forever, depending on the global parameters describing the intensity of interconnection. This provides a way to understand the emergence of persistent activity triggered by weak input signals in large neural networks.	abstract	This article presents a biological neural network model driven by inhomogeneous Poisson processes accounting for the intrinsic randomness of synapses. The main novelty is the introduction of local interactions: each firing neuron triggers an instantaneous increase in electric potential to a fixed number of randomly chosen neurons. We prove that, as the number of neurons approaches infinity, the finite network converges to a nonlinear meanfield process characterised by a jump-type stochastic differential equation. We show that this process displays a phase transition: the activity of a typical neuron in the infinite network either rapidly dies out, or persists forever, depending on the global parameters describing the intensity of interconnection. This provides a way to understand the emergence of persistent activity triggered by weak input signals in large neural networks.		
	versions		versions			