	doc_1		doc_2		decision	id
cases	authors	Manivel, Laurent	authors	Laurent Manivel		
	title	Configurations of lines and models of Lie algebras	title	Configurations of lines and models of Lie algebras		
	publication_date   2005-07-06 00:00:00		publication_date 2005-07-06 00:00:00			
	source	SupportedSources.CORE	source	SupportedSources.INTERNET_ARCHIVE		
	journal		journal			
	volume		volume			
	doi	10.1016/j.jalgebra.2006.04.029	doi			
	urls	http://arxiv.org/abs/math/0507118	urls	https://archive.org/download/arxiv-math0507118/math0507118.pdf		904
	id	id-7919433715303589922	id	id1796320959106989510		
	abstract	The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of \$E\_6\$ and \$E\_7\$. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For \${\mathfrak e}\_7\$ and \${\mathfrak e}\_8\$ we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism \$PSL(3,F\_2)\simeq PSL(2,F\_7)\$ in terms of harmonic cubes.Comment: 31 page	abstract	The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of E_6 and E_7. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For e_7 and e_8 we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism PSL(3,F_2)â‰f PSL(2,F_7) in terms of harmonic cubes.		
	versions		versions			