

cases	doc_1		doc_2		decision	id
					DUPLICATES	379
	authors	<ul style="list-style-type: none">Dorodnyi, M.Suslina, T.	authors	<ul style="list-style-type: none">Mark DorodnyiTatiana Suslina		
	title	Spectral approach to homogenization of hyperbolic equations with periodic coefficients	title	Spectral approach to homogenization of hyperbolic equations with periodic coefficients		
	publication_date	2018-01-01 00:00:00	publication_date	2017-08-02 07:22:23+00:00		
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV		
	journal		journal	None		
	volume		volume			
	doi	10.1016/j.jde.2018.02.023	doi			
	urls	<ul style="list-style-type: none">https://api.elsevier.com/content/article/PII:S0022039618301128?httpAccept=text/xmlhttps://api.elsevier.com/content/article/PII:S0022039618301128?httpAccept=text/plainhttp://dx.doi.org/10.1016/j.jde.2018.02.023	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1708.00859v1http://arxiv.org/abs/1708.00859v1http://arxiv.org/pdf/1708.00859v1		
	id	id4110978329387452802	id	id-3847912587829905321		
	abstract		abstract	In $L_2(\mathbb{R}^d; \mathbb{C}^n)$, we consider selfadjoint strongly elliptic second order differential operators $\{\mathcal{A}_{\varepsilon}\}$ with periodic coefficients depending on $\{\mathbf{x}/\varepsilon\}$, $\varepsilon>0$. We study the behavior of the operators $\cos(\sqrt{\mathcal{A}_{\varepsilon}}\tau)$ and $\mathcal{A}_{\varepsilon}^{-1/2}\sin(\sqrt{\mathcal{A}_{\varepsilon}}\tau)$, $\tau \in \mathbb{R}$, for small ε . Approximations for these operators in the $(H^s \text{ to } L_2)$ -operator norm with a suitable s are obtained. The results are used to study the behavior of the solution $\{\mathbf{v}_{\varepsilon}\}$ of the Cauchy problem for the hyperbolic equation $\partial_{\tau}^2 \mathbf{v}_{\varepsilon} = -\mathcal{A}_{\varepsilon} \mathbf{v}_{\varepsilon} + \mathbf{F}$. General results are applied to the acoustics equation and the system of elasticity theory.		
	versions		versions			