	doc_1		doc_2		decision	id	
			authors	Joaquim Bruna			
	authors	• J. Bruna	title	L^p estimates for Riesz transforms on forms in the Poincare space H^n			
	title	L^p estimates for Riesz transforms on forms in the Poincare space H^n	publication_date	2004-11-25 09:28:36+00:00			
pul	blication_date	2004-11-25 00:00:00	source	SupportedSources.ARXIV			
	source	SupportedSources.SEMANTIC_SCHOLAR	journal	None			
	journal	arXiv: Analysis of PDEs	volume		DUPLICATES 312		
	volume		doi				
cases	doi urls	https://www.semanticscholar.org/paper/e6f1e77723da123580af3f50791715599e3908fb	urls	 http://arxiv.org/pdf/math/0411569v1 http://arxiv.org/abs/math/0411569v1 http://arxiv.org/pdf/math/0411569v1 		312	
	id	id-7641292473930828713	id	id8473789152861044096			
	abstract versions	Using hyperbolic form convolution with doubly isometry-invariant kernels, the explicit expression of the inverse of the de Rham laplacian acting on m-forms in the Poincar\'{e} space is found. Also, by means of some estimates for hyperbolic singular integrals, we obtain L^p-estimates for the Riesz transforms passing from the Laplacian to other covariant derivatives, in a range of p depending on m,n. Finally, using these, it is shown that the Laplacian defines topological isomorphisms in the scale of form Sobolev spaces, for m different from n/2,(n+1)/2,(n-1)/2.		Using hyperbolic form convolution with doubly isometry-invariant kernels, the explicit expression of the inverse of the de Rham laplacian acting on m-forms in the Poincar\'{e} space is found. Also, by means of some estimates for hyperbolic singular integrals, we obtain L^p-estimates for the Riesz transforms passing from the Laplacian to other covariant derivatives, in a range of p depending on m,n. Finally, using these, it is shown that the Laplacian defines topological isomorphisms in the scale of form Sobolev spaces, for m different from n/2,(n+1)/2,(n-1)/2.			
			versions				