

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Stephan I. TzenovRonald C. Davidson	authors	<ul style="list-style-type: none">Stephan I. TzenovRonald C. Davidson	NOT DUPLICATES	377
	title	Hamiltonian Formalism for Solving the Vlasov-Poisson Equations and Its Application to the Coherent Beam-Beam Interaction	title	Hamiltonian Formalism for Solving the Vlasov-Poisson Equations and Its Applications to Periodic Focusing Systems and the Coherent Beam-Beam Interaction		
	publication_date	2001-06-01 13:33:14+00:00	publication_date	2001-10-04 12:51:20+00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.ARXIV		
	journal	Conf.Proc.C0106181:2075-2077,2001	journal	Phys.Rev.ST Accel.Beams 5 (2002) 021001		
	volume		volume			
	doi		doi	10.1103/PhysRevSTAB.5.021001		
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/physics/0106003v1http://arxiv.org/abs/physics/0106003v1http://arxiv.org/pdf/physics/0106003v1	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/physics/0110014v1http://dx.doi.org/10.1103/PhysRevSTAB.5.021001http://arxiv.org/abs/physics/0110014v1http://arxiv.org/pdf/physics/0110014v1		
	id	id5210095978192341359	id	id3170279473022706645		
	abstract	A Hamiltonian approach to the solution of the Vlasov-Poisson equations has been developed. Based on a nonlinear canonical transformation, the rapidly oscillating terms in the original Hamiltonian are transformed away, yielding a new Hamiltonian that contains slowly varying terms only. The formalism has been applied to the coherent beam-beam interaction, and a stationary solution to the transformed Vlasov equation has been obtained.	abstract	A Hamiltonian approach to the solution of the Vlasov-Poisson equations has been developed. Based on a nonlinear canonical transformation, the rapidly oscillating terms in the original Hamiltonian are transformed away, yielding a new Hamiltonian that contains slowly varying terms only. The formalism has been applied to the dynamics of an intense beam propagating through a periodic focusing lattice, and to the coherent beam-beam interaction. A stationary solution to the transformed Vlasov equation has been obtained.		
	versions		versions			