

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">E. KharazmiZ. ZhangG. E. Karniadakis	authors	<ul style="list-style-type: none">E. KharazmiZ. ZhangG. Karniadakis	DUPLICATES	78
	title	Variational Physics-Informed Neural Networks For Solving Partial Differential Equations	title	Variational Physics-Informed Neural Networks For Solving Partial Differential Equations		
	publication_date	2019-11-27 19:51:39+00:00	publication_date	2019-11-27 00:00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	None	journal	ArXiv		
	volume		volume	abs/1912.00873		
	doi		doi			
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1912.00873v1http://arxiv.org/abs/1912.00873v1http://arxiv.org/pdf/1912.00873v1	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/d483f6ecc4685767344822d9e0f03c82b68531ba		
	id	id-8620909391592721593	id	id-3217945672895159295		
	abstract	Physics-informed neural networks (PINNs) [31] use automatic differentiation to solve partial differential equations (PDEs) by penalizing the PDE in the loss function at a random set of points in the domain of interest. Here, we develop a Petrov-Galerkin version of PINNs based on the nonlinear approximation of deep neural networks (DNNs) by selecting the {\em trial space} to be the space of neural networks and the {\em test space} to be the space of Legendre polynomials. We formulate the \textit{variational residual} of the PDE using the DNN approximation by incorporating the variational form of the problem into the loss function of the network and construct a \textit{variational physics-informed neural network} (VPINN). By integrating by parts the integrand in the variational form, we lower the order of the differential operators represented by the neural networks, hence effectively reducing the training cost in VPINNs while increasing their accuracy compared to PINNs that essentially employ delta test functions. For shallow networks with one hidden layer, we analytically obtain explicit forms of the \textit{variational residual}. We demonstrate the performance of the new formulation for several examples that show clear advantages of VPINNs over PINNs in terms of both accuracy and speed.	abstract	Physics-informed neural networks (PINNs) [31] use automatic differentiation to solve partial differential equations (PDEs) by penalizing the PDE in the loss function at a random set of points in the domain of interest. Here, we develop a Petrov-Galerkin version of PINNs based on the nonlinear approximation of deep neural networks (DNNs) by selecting the {\em trial space} to be the space of neural networks and the {\em test space} to be the space of Legendre polynomials. We formulate the \textit{variational residual} of the PDE using the DNN approximation by incorporating the variational form of the problem into the loss function of the network and construct a \textit{variational physics-informed neural network} (VPINN). By integrating by parts the integrand in the variational form, we lower the order of the differential operators represented by the neural networks, hence effectively reducing the training cost in VPINNs while increasing their accuracy compared to PINNs that essentially employ delta test functions. For shallow networks with one hidden layer, we analytically obtain explicit forms of the \textit{variational residual}. We demonstrate the performance of the new formulation for several examples that show clear advantages of VPINNs over PINNs in terms of both accuracy and speed.		
	versions		versions			