

cases	doc_1		doc_2				decision	id
			authors	<ul style="list-style-type: none"><li>Anna V. Anop</li><li>Aleksandr A. Murach</li></ul>			DUPLICATES	1232
	authors	<ul style="list-style-type: none"><li>Anna Anop</li><li>Aleksandr A. Murach</li></ul>	title	Parameter-elliptic problems and interpolation with a function parameter				
	title	Parameter-elliptic problems and interpolation with a function parameter	publication_date	2014-03-11 11:40:36+00:00				
	publication_date	2014-03-11 00:00:00	source	SupportedSources.ARXIV				
	source	SupportedSources.OPENALEX	journal	Methods Funct. Anal. Topology 20 (2014), no. 2, 103-116				
	journal	arXiv (Cornell University)	volume					
	volume		doi					
	doi	None	urls	<ul style="list-style-type: none"><li>http://arxiv.org/pdf/1403.2542v1</li><li>http://arxiv.org/abs/1403.2542v1</li><li>http://arxiv.org/pdf/1403.2542v1</li></ul>				
	urls	<ul style="list-style-type: none"><li>https://openalex.org/W1523053138</li></ul>	id	id-1330728505637596651				
	id	id-793696308483509573	abstract	Parameter-elliptic boundary-value problems are investigated on the extended Sobolev scale. This scale consists of all Hilbert spaces that are interpolation spaces with respect to the Hilbert Sobolev scale. The latter are the $H^s$ -ormander spaces $B_{2,k}^s$ for which the smoothness index $s$ is an arbitrary radial function $\rho$ -varying at infinity. We prove that the operator corresponding to this problem sets isomorphisms between appropriate $H^s$ -ormander spaces provided that the absolute value of the parameter is large enough. For solutions to the problem, we establish two-sided estimates, in which the constants are independent of the parameter.				
	abstract		versions					
	versions							