

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>Anna Anop</li><li>Aleksandr A. Murach</li></ul>	authors	<ul style="list-style-type: none"><li>A. Anop</li><li>A. Murach</li></ul>	DUPLICATES	1231
	title	Parameter-elliptic problems and interpolation with a function parameter	title	Parameter-elliptic problems and interpolation with a function parameter		
	publication_date	2014-03-11 00:00:00	publication_date	2014-03-11 00:00:00		
	source	SupportedSources.OPENALEX	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	arXiv (Cornell University)	journal	arXiv: Analysis of PDEs		
	volume		volume			
	doi	None	doi			
	urls	<ul style="list-style-type: none"><li>https://openalex.org/W1523053138</li></ul>	urls	<ul style="list-style-type: none"><li>https://www.semanticscholar.org/paper/b23afae1b399f6620698b29f837c5202f1500e7d</li></ul>		
	id	id-793696308483509573	id	id4737350623277622325		
	abstract		abstract	Parameter-elliptic boundary-value problems are investigated on the extended Sobolev scale. This scale consists of all Hilbert spaces that are interpolation spaces with respect to the Hilbert Sobolev scale. The latter are the $H^s$ -ormander spaces $B_{2,k}^s$ for which the smoothness index $s$ is an arbitrary radial function $\rho$ -varying at infinity. We prove that the operator corresponding to this problem sets isomorphisms between appropriate $H^s$ -ormander spaces provided that the absolute value of the parameter is large enough. For solutions to the problem, we establish two-sided estimates, in which the constants are independent of the parameter.		
	versions		versions			