cases	doc_1	doc_2	decision	id
	<table><tr><td>**authors**</td><td>• Anna Anop • Aleksandr A. Murach</td></tr><tr><td>**title**</td><td>Parameter-elliptic problems and interpolation with a function parameter</td></tr><tr><td>**publication_date**</td><td>2014-03-11 00:00:00</td></tr><tr><td>**source**</td><td>SupportedSources.OPENALEX</td></tr><tr><td>**journal**</td><td>arXiv (Cornell University)</td></tr><tr><td>**volume**</td><td></td></tr><tr><td>**doi**</td><td>None</td></tr><tr><td>**urls**</td><td>• https://openalex.org/W1523053138</td></tr><tr><td>**id**</td><td>id-793696308483509573</td></tr><tr><td>**abstract**</td><td></td></tr><tr><td>**versions**</td><td></td></tr></table>	<table><tr><td>**authors**</td><td>• Anna V. Anop • Aleksandr A. Murach</td></tr><tr><td>**title**</td><td>Parameter-elliptic problems and interpolation with a function parameter</td></tr><tr><td>**publication_date**</td><td>2014-03-11 00:00:00</td></tr><tr><td>**source**</td><td>SupportedSources.INTERNET_ARCHIVE</td></tr><tr><td>**journal**</td><td></td></tr><tr><td>**volume**</td><td></td></tr><tr><td>**doi**</td><td></td></tr><tr><td>**urls**</td><td>• https://web.archive.org/web/20191022101140/https://arxiv.org/pdf/1403.2542v1.pdf</td></tr><tr><td>**id**</td><td>id7176659057869114802</td></tr><tr><td>**abstract**</td><td>Parameter-elliptic boundary-value problems are investigated on the extended Sobolev scale. This scale consists of all Hilbert spaces that are interpolation spaces with respect to the Hilbert Sobolev scale. The latter are the HÃ¶rmander spaces B_2,k for which the smoothness index k is an arbitrary radial function RO-varying at infinity. We prove that the operator corresponding to this problem sets isomorphisms between appropriate HÃ¶rmander spaces provided that the absolute value of the parameter is large enough. For solutions to the problem, we establish two-sided estimates, in which the constants are independent of the parameter.</td></tr><tr><td>**versions**</td><td></td></tr></table>	DUPLICATES	1233