

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>Mazen Ali</li><li>Anthony Nouy</li></ul>	authors	<ul style="list-style-type: none"><li>Mazen Ali</li><li>Anthony Nouy</li></ul>	DUPLICATES	98
	title	Singular Value Decomposition in Sobolev Spaces: Part I	title	Singular Value Decomposition in Sobolev Spaces: Part I		
	publication_date	2019-12-25 00:00:00	publication_date	2019-12-25 00:00:00		
	source	SupportedSources.INTERNET_ARCHIVE	source	SupportedSources.INTERNET_ARCHIVE		
	journal		journal			
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none"><li>https://web.archive.org/web/20200321104102/https://arxiv.org/pdf/1809.11001v3.pdf</li></ul>	urls	<ul style="list-style-type: none"><li>https://web.archive.org/web/20200321104102/https://arxiv.org/pdf/1809.11001v3.pdf</li></ul>		
	id	id8511692479879353716	id	id5256768460988041481		
	abstract	A well known result from functional analysis states that any compact operator between Hilbert spaces admits a singular value decomposition (SVD). This decomposition is a powerful tool that is the workhorse of many methods both in mathematics and applied fields. A prominent application in recent years is the approximation of high-dimensional functions in a low-rank format. This is based on the fact that, under certain conditions, a tensor can be identified with a compact operator and SVD applies to the latter. One key assumption for this application is that the tensor product norm is not weaker than the injective norm. This assumption is not fulfilled in Sobolev spaces, which are widely used in the theory and numerics of partial differential equations. Our goal is the analysis of the SVD in Sobolev spaces. This work consists of two parts. In this manuscript (part I), we address low-rank approximations and minimal subspaces in H1. We analyze the H1-error of the SVD performed in the ambient L2-space. In part II, we will address variants of the SVD in norms stronger than the L2-norm. We will provide a few numerical examples that support our theoretical findings.	abstract	A well known result from functional analysis states that any compact operator between Hilbert spaces admits a singular value decomposition (SVD). This decomposition is a powerful tool that is the workhorse of many methods both in mathematics and applied fields. A prominent application in recent years is the approximation of high-dimensional functions in a low-rank format. This is based on the fact that, under certain conditions, a tensor can be identified with a compact operator and SVD applies to the latter. One key assumption for this application is that the tensor product norm is not weaker than the injective norm. This assumption is not fulfilled in Sobolev spaces, which are widely used in the theory and numerics of partial differential equations. Our goal is the analysis of the SVD in Sobolev spaces. This work consists of two parts. In this manuscript (part I), we address low-rank approximations and minimal subspaces in H1. We analyze the H1-error of the SVD performed in the ambient L2-space. In part II, we will address variants of the SVD in norms stronger than the L2-norm. We will provide a few numerical examples that support our theoretical findings.		
	versions		versions			