	doc_1		doc_2		decision	id
cases	authors	Ian Holloway Sivaguru S. Sritharan				
	title	Ideal Magnetohydrodynamic Equations on a Sphere and Elliptic-Hyperbolic Property	authors	Ian Holloway S. Sritharan		
		cation date 2019-10-20 04:28:58+00:00		5. Sitthurun		
		SupportedSources.ARXIV	title	Ideal magnetohydrodynamic equations on a sphere and elliptic-hyperbolic property		
	source	None Supported Sources. ARXIV	publication_date 2019-10-20 00:00:00			
	journal volume	None	source	SupportedSources.SEMANTIC_SCHOLAR		
			journal			
	doi	1447/17 2 201/21/1010 10/27 1	volume			
	urls	 http://arxiv.org/pdf/1910.10637v1 http://arxiv.org/abs/1910.10637v1 http://arxiv.org/pdf/1910.10637v1 	doi	10.1090/qam/1571		
			urls	https://www.semanticscholar.org/paper/83dd74a5a727b816362d275e9bf8908aedb7a42d		1019
	id	id3814579275512915648	id	id-812290558807850697		
	abstract	This work contains the derivation and type analysis of the conical Ideal Magnetohydrodynamic equations. The 3D Ideal MHD equations with Powell source terms, subject to the assumption that the solution is conically invariant, are projected onto a unit sphere using tools from tensor calculus. Conical flows provide valuable insight into supersonic and hypersonic flow past bodies, but are simpler to analyze and solve numerically. Previously, work has been done on conical inviscid flows governed by the Euler equations with great success. It is known that some flight regimes involve flows of ionized gases, and thus there is motivation to extend the study of conical flows to the case where the gas is electrically conducting. To the authors' knowledge, the conical magnetohydrodynamic equations have never been derived and so this paper is the first invesitgation of that system. Among the results, we show that conical flows for this case do exist mathematically and that the governing system of partial differential equations is of mixed type. Throughout the domain it	abstract versions	This work contains the derivation and type analysis of the conical ideal magnetohydrodynamic equations. The 3D ideal MHD equations with Powell source terms, subject to the assumption that the solution is conically invariant, are projected onto a unit sphere using tools from tensor calculus. Conical flows provide valuable insight into supersonic and hypersonic flow past bodies, but are simpler to analyze and solve numerically. Previously, work has been done on conical inviscid flows governed by the Euler equations with great success. It is known that some flight regimes involve flows of ionized gases, and thus there is motivation to extend the study of conical flows to the case where the gas is electrically conducting. To the authors' knowledge, the conical magnetohydrodynamic equations have never been derived, and so this paper is the first investigation of that system. Among the results, we show that conical flows for this case do exist mathematically and that the governing system of partial differential equations is of mixed type. Throughout the domain it can be either hyperbolic or elliptic depending on the solution.		
	versions	can be either hyperbolic or elliptic depending on the solution.				