	doc_1		doc_2		decision	id
	authors	A. Ardjouni A. Djoudi	authors	A. Ardjouni A. Djoudi		
	title	The existence of periodic solutions for a second order nonlinear neutral differential equation with functional delay	title	EXISTENCE OF PERIODIC SOLUTIONS FOR A SECOND ORDER NONLINEAR NEUTRAL FUNCTIONAL DIFFERENTIAL EQUATION		
	publication_date None		publication_date None			
	source	SupportedSources.SEMANTIC_SCHOLAR	source	SupportedSources.SEMANTIC_SCHOLAR		1
	journal	Electronic Journal of Qualitative Theory of Differential Equations	journal			
cases	volume		volume		NOT	
	doi	10.14232/EJQTDE.2012.1.31	doi		DUPLICATES	531
	urls	https://www.semanticscholar.org/paper/9e0d3fa3db76e1d2d7e2a0ecff4e61eabb46ae43	urls	https://www.semanticscholar.org/paper/bc9e2251fff8e79e07ee3e110f884c4ea8ce96e8		
	id	id-1832595271075805063	id	id-7486091728476523865		
	abstract	In this article we study the existence of periodic solutions of the second order nonlinear neutral differential equation with functional delay d dt x (t) + p (t) d dt x (t) + q (t) x (t) = d dt g (t, x (tâ^' I , (t))) + f ' t, x (t), x (tâ^' I , (t)) I . The main tool employed here is the Burton-Krasnoselskiiâ E^{TM} s hybrid fixed point theorem dealing with a sum of two mappings, one is a large contraction and the other is compact.	abstract	We study the existence of periodic solutions of the second order nonlinear neutral differential equation with variable delay $x \hat{a} \in \hat{a} \in \hat{c}$ (t) + p (t) $x \hat{a} \in \hat{c}$ (t) + q (t)h (x (t)) = c (t) $x \hat{a} \in \hat{c}$ (t\hat{a}^* \tilde{I}, (t)) + f (t, x (t\hat{a}^* \tilde{I}, (t))). We invert the given equation to obtain an integral, but equivalent, equation from which we define a fixed point mapping written as a sum of a large contraction and a compact map. We show that such maps fit very nicely into the framework of Krasnoselskii-Burton\hat{a} \in \text{TMs} fixed point theorem so that the existence of periodic solutions is conclued.		
	versions		versions			'