

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"> Johannes Huebschmann 	authors	<ul style="list-style-type: none"> Johannes Huebschmann 	NOT DUPLICATES	1248
	title	Smooth structures on certain moduli spaces for bundles on a surface	title	Poisson structures on certain moduli spaces for bundles on a surface		
	publication_date	1994-11-23 09:51:18+00:00	publication_date	1994-11-23 10:03:51+00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.ARXIV		
	journal	J. of Pure and Applied Algebra 128 (1998), 183-221	journal	Ann. Inst. Fourier 45 (1995) 65-91		
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none"> http://arxiv.org/pdf/dg-ga/9411008v1 http://arxiv.org/abs/dg-ga/9411008v1 http://arxiv.org/pdf/dg-ga/9411008v1 	urls	<ul style="list-style-type: none"> http://arxiv.org/pdf/dg-ga/9411009v1 http://arxiv.org/abs/dg-ga/9411009v1 http://arxiv.org/pdf/dg-ga/9411009v1 		
	id	id-7604876183244006927	id	id6540082670780602496		
	abstract	Let Σ be a closed surface, G a compact Lie group, with Lie algebra \mathfrak{g} , $\mathcal{P} \rightarrow \Sigma$ a principal G -bundle, let \mathcal{N} denote the moduli space of central Yang-Mills connections on Σ , for suitably chosen additional data, and let $\text{Rep}_\chi(\Gamma, G)$ be the space of representations of the universal central extension $\tilde{\Gamma}$ of Γ in G that corresponds to χ . We construct smooth structures on \mathcal{N} and $\text{Rep}_\chi(\Gamma, G)$ and show that the assignment to a smooth connection A of its holonomies with reference to suitable closed paths yields a diffeomorphism from \mathcal{N} onto $\text{Rep}_\chi(\Gamma, G)$; moreover we show that the derivative of the latter at the non-singular points of \mathcal{N} amounts to a certain twisted integration mapping. Finally we examine the infinitesimal geometry of these moduli spaces with reference to the smooth structures and, for illustration, we show that, on the moduli space of flat $\text{SU}(2)$ -connections for a surface of genus two which, as a space, is just complex projective 3-space, our smooth structure looks rather different from the standard structure.	abstract	Let Σ be a closed surface, G a compact Lie group, with Lie algebra \mathfrak{g} , and $\mathcal{P} \rightarrow \Sigma$ a principal G -bundle. In earlier work we have shown that the moduli space \mathcal{N} of central Yang-Mills connections, for appropriate additional data, is stratified by smooth symplectic manifolds and that the holonomy yields a diffeomorphism from \mathcal{N} onto a certain representation space $\text{Rep}_\chi(\Gamma, G)$, with reference to suitable smooth structures $C^\infty(\mathcal{N})$ and $C^\infty(\text{Rep}_\chi(\Gamma, G))$ where $\tilde{\Gamma}$ denotes the universal central extension of the fundamental group of Σ . Given an invariant symmetric bilinear form on \mathfrak{g}^* , we construct here Poisson structures on $C^\infty(\mathcal{N})$ and $C^\infty(\text{Rep}_\chi(\Gamma, G))$ in such a way that the mentioned diffeomorphism identifies them. When the form on \mathfrak{g}^* is non-degenerate the Poisson structures are compatible with the stratifications where $\text{Rep}_\chi(\Gamma, G)$ is endowed with the corresponding stratification and, furthermore, yield structures of a stratified symplectic space, preserved by the induced action of the mapping class group of Σ .		
	versions		versions			