

cases	doc_1		doc_2		decision	id
			authors	<ul style="list-style-type: none">Adam, John A.	NOT DUPLICATES	1179
	authors	<ul style="list-style-type: none">John A. Adam	title	A Nonlinear Eigenvalue Problem in Astrophysical Magnetohydrodynamics: Some Properties of the Spectrum		
	title	A nonlinear eigenvalue problem in astrophysical magnetohydrodynamics: Some properties of the spectrum	publication_date	1989-01-01 00:00:00		
	publication_date	1989-03-01 00:00:00	source	SupportedSources.CORE		
	source	SupportedSources.OPENALEX	journal			
	journal	Journal of Mathematical Physics	volume			
	volume		doi	None		
	doi	10.1063/1.528392	urls	<ul style="list-style-type: none">https://core.ac.uk/download/217299455.pdf		
	urls	<ul style="list-style-type: none">https://openalex.org/W2046262924https://doi.org/10.1063/1.528392https://digitalcommons.odu.edu/cgi/viewcontent.cgi?article=1153&context=mathstat_fac_pubs	id	id-5345294098406050429		
	id	id8680548999588315659	abstract	The equations of ideal magnetohydrodynamics (MHD) with an external gravitational potential $\phi_a = \hat{r}^{\alpha}$ "magnetoatmosphere" $r^{TM\alpha TM\beta}$ are examined in detail as a singular nonlinear eigenvalue problem. Properties of the spectrum are discussed with specific emphasis on two systems relevant to solar magnetohydrodynamics. In the absence of a gravitational potential, the system reduces to that of importance in MHD and plasma physics, albeit in a different geometry. This further reduces to a form isomorphic to that derived in the study of plasma oscillations in a cold plasma, Alfvén wave propagation in an inhomogeneous medium, and MHD waves in a sheet pinch. In cylindrical geometry, the relevant model equations are those for a diffuse linear pinch. The full system, including gravity, has been applied to the study of flare-induced coronal waves, running penumbral waves in sunspots, and linear wave coupling in a highly inhomogeneous medium. The structure of the so-called MHD critical layer and its contribution to the continuous spectrum is examined in detail for a model magnetoatmosphere, based on properties of the hypergeometric differential operator. The relationship of this singular region to critical layers in classical linear hydrodynamic stability theory is also discussed in the light of a specific model (in the Appendix)		
	abstract		versions			
	versions					