	doc_1		doc_2		decision	id
cases	authors	L. Manivel	authors	Laurent Manivel		
	title	Configurations of lines and models of Lie algebras	title	Configurations of lines and models of Lie algebras	s, ad	
		2006-01-01 00:00:00	<del></del>	2005-07-06 00:00:00 SupportedSources.INTERNET ARCHIVE		
		SupportedSources.INTERNET_ARCHIVE	journal	Supported Sources. IN TERNET_TREE IT		
	<b>J</b>	Elsevier BV	volume			
	volume doi	10.1016/i iologbys 2006.04.020	doi			
	urls	10.1016/j.jalgebra.2006.04.029   https://web.archive.org/web/20190417015936/https://core.ac.uk/download/pdf/81999529.pdf	urls	https://archive.org/download/arxiv-math0507118/math0507118.pdf		898
	id	id-9200781509194487460	id	id1796320959106989510		
	abstract	The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of E 6 and E 7. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For e 7 and e 8 we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism PSL(3, F 2) PSL(2, F 7) in terms of harmonic cubes.		The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of E_6 and E_7. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For e_7 and e_8 we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism PSL(3,F_2)â‰f PSL(2,F_7) in terms of harmonic cubes.		
	versions		versions			