

| cases | doc_1            |   | doc_2            |   | decision   | id |
|-------|------------------|---|------------------|---|------------|----|
|       |                  |   |                  |   | DUPLICATES | 86 |
|       | authors          | <ul style="list-style-type: none"><li>M. Dorodnyi</li><li>T. Suslina</li></ul>  | authors          | <ul style="list-style-type: none"><li>Mark Dorodnyi</li><li>Tatiana Suslina</li></ul>   |            |    |
|       | title            | Homogenization of hyperbolic equations with periodic coefficients   | title            | Homogenization of hyperbolic equations with periodic coefficients   |            |    |
|       | publication_date | 2016-06-19 00:00:00   | publication_date | 2016-06-19 13:27:50+00:00   |            |    |
|       | source           | SupportedSources.SEMANTIC_SCHOLAR   | source           | SupportedSources.ARXIV  |            |    |
|       | journal          | arXiv: Analysis of PDEs   | journal          | None  |            |    |
|       | volume           |   | volume           |   |            |    |
|       | doi              |   | doi              |   |            |    |
|       | urls             | <ul style="list-style-type: none"><li>https://www.semanticscholar.org/paper/7aa66b7786bee542c3afed84f2c55fbc00956241</li></ul>  | urls             | <ul style="list-style-type: none"><li>http://arxiv.org/pdf/1606.05868v1</li><li>http://arxiv.org/abs/1606.05868v1</li><li>http://arxiv.org/pdf/1606.05868v1</li></ul>   |            |    |
|       | id               | id2794005982141959912   | id               | id6618946359099955088   |            |    |
|       | abstract         | In $L_2(\mathbb{R}^d; \mathbb{C}^n)$ we consider selfadjoint strongly elliptic second order differential operators $\mathcal{A}_\varepsilon$ with periodic coefficients depending on $\mathbf{x}/\varepsilon$ , $\varepsilon>0$ . We study the behavior of the operator cosine $\cos(\sqrt{\mathcal{A}_\varepsilon}/2\varepsilon)\tau$ , $\tau\in\mathbb{R}$ , for small $\varepsilon$ . Approximations for this operator in the $(H^s\text{to }L_2)$ -operator norm with a suitable $s$ are obtained. The results are used to study the behavior of the solution $\mathbf{v}_\varepsilon$ of the Cauchy problem for the hyperbolic equation $\partial_\tau^2 \mathbf{v}_\varepsilon = -\mathcal{A}_\varepsilon \mathbf{v}_\varepsilon + \mathbf{F}$ . General results are applied to the acoustics equation and the system of elasticity theory. | abstract         | In $L_2(\mathbb{R}^d; \mathbb{C}^n)$ we consider selfadjoint strongly elliptic second order differential operators $\mathcal{A}_\varepsilon$ with periodic coefficients depending on $\mathbf{x}/\varepsilon$ , $\varepsilon>0$ . We study the behavior of the operator cosine $\cos(\sqrt{\mathcal{A}_\varepsilon}/2\varepsilon)\tau$ , $\tau\in\mathbb{R}$ , for small $\varepsilon$ . Approximations for this operator in the $(H^s\text{to }L_2)$ -operator norm with a suitable $s$ are obtained. The results are used to study the behavior of the solution $\mathbf{v}_\varepsilon$ of the Cauchy problem for the hyperbolic equation $\partial_\tau^2 \mathbf{v}_\varepsilon = -\mathcal{A}_\varepsilon \mathbf{v}_\varepsilon + \mathbf{F}$ . General results are applied to the acoustics equation and the system of elasticity theory. |            |    |
|       | versions         |   | versions         |   |            |    |