

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Bin ChengQiangchang JuSteve Schochet	authors	<ul style="list-style-type: none">B. ChengQiangchang JuS. Schochet	DUPLICATES	1014
	title	Convergence Rate Estimates for the Low Mach and Alfvén Number Three-Scale Singular Limit of Compressible Ideal Magnetohydrodynamics	title	Convergence rate estimates for the low Mach and Alfvén number three-scale singular limit of compressible ideal magnetohydrodynamics		
	publication_date	2019-03-14 04:58:13+00:00	publication_date	2019-03-14 00:00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	None	journal			
	volume		volume			
	doi		doi	10.1051/m2an/2020051		
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1903.05816v3http://arxiv.org/abs/1903.05816v3http://arxiv.org/pdf/1903.05816v3	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/1e2912ec5a35faa4e561f44c47771af7f94d7cf4		
	id	id7860676359247038454	id	id-8673461365415629256		
	abstract	Convergence rate estimates are obtained for singular limits of the compressible ideal magnetohydrodynamics equations, in which the Mach and Alfvén numbers tend to zero at different rates. The proofs use a detailed analysis of exact and approximate fast, intermediate, and slow modes together with improved estimates for the solutions and their time derivatives, and the time-integration method. When the small parameters are related by a power law the convergence rates are positive powers of the Mach number, with the power varying depending on the component and the norm. Exceptionally, the convergence rate for two components involve the ratio of the two parameters, and that rate is proven to be sharp via corrector terms. Moreover, the convergence rates for the case of a power-law relation between the small parameters tend to the two-scale convergence rate as the power tends to one. These results demonstrate that the issue of convergence rates for three-scale singular limits, which was not addressed in the authors' previous paper, is much more complicated than for the classical two-scale singular limits.	abstract	Convergence rate estimates are obtained for singular limits of the compressible ideal magnetohydrodynamics equations, in which the Mach and Alfvén numbers tend to zero at different rates. The proofs use a detailed analysis of exact and approximate fast, intermediate, and slow modes together with improved estimates for the solutions and their time derivatives, and the time-integration method. When the small parameters are related by a power law the convergence rates are positive powers of the Mach number, with the power varying depending on the component and the norm. Exceptionally, the convergence rate for two components involve the ratio of the two parameters, and that rate is proven to be sharp via corrector terms. Moreover, the convergence rates for the case of a power-law relation between the small parameters tend to the two-scale convergence rate as the power tends to one. These results demonstrate that the issue of convergence rates for three-scale singular limits, which was not addressed in the authors' previous paper, is much more complicated than for the classical two-scale singular limits.		
	versions		versions			