

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>K. Xu</li></ul>	authors	<ul style="list-style-type: none"><li>Kun Xu Icase</li></ul>	DUPLICATES	30
	title	Gas-Kinetic Theory-Based Flux Splitting Method for Ideal Magnetohydrodynamics	title	Gas-Kinetic Theory-Based Flux Splitting Method for Ideal Magnetohydrodynamics		
	publication_date	1999-08-10 00:00:00	publication_date	None		
	source	SupportedSources.SEMANTIC_SCHOLAR	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	Journal of Computational Physics	journal			
	volume	153	volume			
	doi	10.1006/JCPH.1999.6280	doi			
	urls	<ul style="list-style-type: none"><li>https://www.semanticscholar.org/paper/d54b16a1164984c383bdd5978abd861b1887dd82</li></ul>	urls	<ul style="list-style-type: none"><li>https://www.semanticscholar.org/paper/80915df07e2931ee9af2b2dfa345c7b0789f7c16</li></ul>		
	id	id-5741628443351207850	id	id-2977725482136893785		
		<p>A gas-kinetic flux splitting method is developed for the ideal magnetohydrodynamics (MHD) equations. The new scheme is based on the direct splitting of the flux function of the MHD equations with the inclusion of “particle” collisions in the transport process. Consequently, the artificial dissipation in the new scheme is greatly reduced in comparison with the MHD flux vector splitting method. Numerical results from the current scheme are favorable compared with those from the well-developed Roe-type MHD solver. In the current paper, the general principle of splitting the macroscopic flux function based on the gas-kinetic theory is presented. The flux construction strategy may shed some light on the possible construction of accurate and robust hybrid schemes for the compressible flow simulations.</p>			<p>A gas-kinetic flux splitting method is developed for the ideal magnetohydrodynamics (MHD) equations. The new scheme is based on the direct splitting of the flux function of the MHD equations with the inclusion of “particle” collisions in the transport process. Consequently, the artificial dissipation in the new scheme is greatly reduced in comparison with the MHD flux vector splitting method. Numerical results from the current scheme are favorable compared with those from the well-developed Roe-type MHD solver. In the current paper, the general principle of splitting the macroscopic flux function based on the gas-kinetic theory is presented. The flux construction strategy may shed some light on the possible construction of accurate and robust hybrid schemes for the compressible flow simulations. c © 1999 Academic Press</p>	
		versions				