

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>Xumin Gu</li></ul>			NOT DUPLICATES	1239
	title	Well-posedness of axially symmetric incompressible ideal magnetohydrodynamic equations with vacuum under the Rayleigh-Taylor sign condition	authors	<ul style="list-style-type: none"><li>Xumin Gu</li></ul>		
	publication_date	2017-12-06 12:14:33+00:00	title	Well-posedness of axially symmetric incompressible ideal magnetohydrodynamic equations with vacuum under the non-collinearity condition		
	source	SupportedSources.ARXIV	publication_date	2017-11-23 00:00:00		
	journal	None	source	SupportedSources.INTERNET_ARCHIVE		
	volume		journal			
	doi		volume			
	urls	<ul style="list-style-type: none"><li>http://arxiv.org/pdf/1712.02152v1</li><li>http://arxiv.org/abs/1712.02152v1</li><li>http://arxiv.org/pdf/1712.02152v1</li></ul>	doi			
	id	id-7065163989295393136	urls	<ul style="list-style-type: none"><li>https://web.archive.org/web/20191017135143/https://arxiv.org/pdf/1711.09757v1.pdf</li></ul>		
	abstract	We consider a free boundary problem for the axially symmetric incompressible ideal magnetohydrodynamic equations that describes the motion of the plasma in vacuum. Both the plasma magnetic field and vacuum magnetic field are tangent along the plasma-vacuum interface. Moreover, the vacuum magnetic field is composed in a non-simply connected domain and hence is non-trivial. Under the Rayleigh-Taylor sign condition on the free surface, we prove the local well-posedness of the problem in Sobolev spaces. Furthermore, we also prove the local well-posdeness under a more general "stability" assumption for the initial data, which provided that the Rayleigh-Taylor sign condition is satisfied at all those points of the initial interface where the non-collinearity condition fails.	id	id-6052351515676971318		
	versions		abstract	We consider a free boundary problem for the axially symmetric incompressible ideal magnetohydrodynamic equations that describes the motion of the plasma in vacuum. Both the plasma magnetic field and vacuum magnetic field are tangent along the plasma-vacuum interface. Moreover, the vacuum magnetic field is composed in a non-simply connected domain and hence is non-trivial. Under the non-collinearity condition on the free surface, we prove the local well-posedness of the problem in Sobolev spaces.		
			versions			