

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Nguyen LamGuozhen Lu	authors	<ul style="list-style-type: none">Nguyen LamGuozhen Lu	NOT DUPLICATES	1932
	title	Sharp singular Adams inequalities in high order Sobolev spaces	title	Sharp Adams type inequalities in Sobolev spaces $W^{m,\frac{n}{m}}(\mathbb{R}^n)$ for arbitrary integer m		
	publication_date	2011-12-29 20:58:09+00:00	publication_date	2011-12-29 19:48:08+00:00		
	source	SupportedSources.ARXIV	source	SupportedSources.ARXIV		
	journal	None	journal	None		
	volume		volume			
	doi		doi			
	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1112.6431v1http://arxiv.org/abs/1112.6431v1http://arxiv.org/pdf/1112.6431v1	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1112.6397v1http://arxiv.org/abs/1112.6397v1http://arxiv.org/pdf/1112.6397v1		
	id	id2785754343849818854	id	id-7143352130776687496		
	abstract	In this paper, we prove a version of weighted inequalities of exponential type for fractional integrals with sharp constants in any domain of finite measure in \mathbb{R}^n . Using this we prove a sharp singular Adams inequality in high order Sobolev spaces in bounded domain at critical case. Then we prove sharp singular Adams inequalities for high order derivatives on unbounded domains. Our results extend the singular Moser-Trudinger inequalities of first order in \cite{Ad2, R, LR, AdY} to the higher order Sobolev spaces $W^{m,\frac{n}{m}}$ and the results of \cite{RS} on Adams type inequalities in unbounded domains to singular case. Our singular Adams inequality on $W^{2,2}(\mathbb{R}^4)$ with standard Sobolev norm at the critical case settles a unsolved question remained in \cite{Y}.	abstract	The main purpose of our paper is to prove sharp Adams-type inequalities in unbounded domains of \mathbb{R}^n for the Sobolev space $W^{m,\frac{n}{m}}(\mathbb{R}^n)$ for any positive integer m less than n . Our results complement those of Ruf and Sani \cite{RS} where such inequalities are only established for even integer m . Our inequalities are also a generalization of the Adams-type inequalities in the special case $n=2m=4$ proved in \cite{Y} and stronger than those in \cite{RS} when $n=2m$ for all positive integer m by using different Sobolev norms.		
	versions		versions			