	doc_1		doc_2		decision	id
	authors	Biegert, M.				
	title	section in the section of the sectio	authors	Markus Biegert		
	publication_date   2009-06-16 00:00:00		title	Lattice Homomorphisms between Sobolev Spaces		
	source	SupportedSources.CROSSREF	rtedSources.CROSSREF publication date 2008-05-30 16:16:21+00:00			
	journal		source	SupportedSources.ARXIV	NOT DUPLICATES 1462	
	volume		journal	None		
cases	doi	10.1007/s11117-009-0022-7	volume			
	urls	<ul> <li>http://link.springer.com/content/pdf/10.1007/s11117- 009-0022-7.pdf</li> <li>http://link.springer.com/article/10.1007/s11117-009- 0022-7/fulltext.html</li> <li>http://link.springer.com/content/pdf/10.1007/s11117- 009-0022-7</li> <li>http://dx.doi.org/10.1007/s11117-009-0022-7</li> <li>id</li> <li>abstract</li> </ul>	doi	• http://arxiv.org/pdf/0805.4740v4		1462
				<ul> <li>http://arxiv.org/abs/0805.4740v4</li> <li>http://arxiv.org/pdf/0805.4740v4</li> </ul>		
			id	id6536245698394927741		
			We show that every vector lattice homomorphism $T$ between Sobolev spaces can be represented by a composition and a multiplication, that is, $T$ is of the form $Tu(x)=u(h(x))g(x)$ for quasi every/almost every $x$ and all $u$ .			
	id	id3976573950376003810	versions	or me term \$ 2 a(m) a(m(m))B(m)\$ ter quarrerery and an \$ aab.		
	abstract		, , , , , , , , , , , , , , , , , , , ,			
	versions					