

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Arielle Leitner	<div></div>		NOT DUPLICATES	1120
	title	A classification of subgroups of $SL(4, \mathbb{R})$ isomorphic to R^3 and generalized cusps in projective 3 manifolds				
	publication_date	2016-01-01 00:00:00				
	source	SupportedSources.INTERNET_ARCHIVE				
	journal	Elsevier BV				
	volume					
	doi	10.1016/j.topol.2016.04.005				
	urls	<ul style="list-style-type: none">https://web.archive.org/web/20180724091901/https://manuscript.elsevier.com/S0166864116300323/pdf/S0166864116300323.pdf				
	id	id-3121647315358814990				
	abstract	<p>This paper uses work of Haettel to classify all subgroups of $PGL(4, \mathbb{R})$ isomorphic to $(R^3, +)$, up to conjugacy. We use this to show there are 4 families of generalized cusps up to projective equivalence in dimension 3. There are 15 conjugacy classes of subgroups in $PGL(4, \mathbb{R})$ isomorphic to $(R^3, +)$, see Theorem 1.1 for the precise list. Classification of abelian subalgebras of $sl(n, \mathbb{R})$ has long been of interest and closely related problems have been studied in [7], [9], [13], and [14]. The remainder of the paper applies this result to classify generalized cusps in convex projective manifolds of dimension 5 3. Suppose M is a manifold of dimension greater than 2. By Mostow-Prasad rigidity, a finite volume hyperbolic structure on M is unique up to isometry. The notion of a hyperbolic structure on a manifold may be generalized to a properly convex projective structure as follows. A convex set with non-empty interior $10 \hat{\mathbb{C}} \hat{\mathbb{S}}, RP^n$ is properly convex if the closure is disjoint from some projective hyperplane. A properly convex manifold is $M = \hat{\mathbb{C}}/\hat{\Gamma}$ where $\hat{\Gamma} \hat{\mathbb{S}}, PGL_{n+1}(\mathbb{R})$ is discrete and acts freely on $\hat{\mathbb{C}}$. If $n = 3$ such an M is a generalized cusp if M is diffeomorphic to $T^2 \hat{\mathbb{A}} \times [0, \hat{a}^* \hat{z})$ and \hat{a}^*, M is strictly convex (contains no line segment). 15 In the case of convex projective structures, there is no notion of Mostow rigidity, so there is a richer deformation theory. When M is closed, Koszul [10], shows that small deformations in the holonomy of hyperbolic structures yield properly convex projective structures. When M is not compact, Koszul's result no longer holds. Cooper, Long and Tillmann [3], have shown Koszul's result 20</p>				
	versions					
	authors	<ul style="list-style-type: none">Arielle Leitner				
	title	A Classification of subgroups of $SL(4, \mathbb{R})$ Isomorphic to R^3 and Generalized Cusps in Projective 3 Manifolds				
	publication_date	2016-04-01 00:00:00				
	source	SupportedSources.INTERNET_ARCHIVE				
	journal					
	volume					
	doi					
	urls	<ul style="list-style-type: none">https://web.archive.org/web/20191027143525/https://arxiv.org/pdf/1507.04724v2.pdf				
	id	id4859770332459554518				
	abstract	This paper uses work of Haettel to classify all subgroups of $PGL(4, \mathbb{R})$ isomorphic to $(R^3, +)$, up to conjugacy. We use this to show there are 4 families of generalized cusps up to projective equivalence in dimension 3.				
	versions					